



Contents lists available at ScienceDirect

Physica A

journal homepage: [www.elsevier.com/locate/physa](http://www.elsevier.com/locate/physa)

# On spin systems with quenched randomness: Classical and quantum

Rafael L. Greenblatt<sup>a,\*</sup>, Michael Aizenman<sup>b,c</sup>, Joel L. Lebowitz<sup>a,d</sup>

<sup>a</sup> Department of Physics and Astronomy, Rutgers University, Piscataway NJ 08854-8019, USA

<sup>b</sup> Department of Physics, Princeton University, Princeton NJ 08544, USA

<sup>c</sup> Department of Mathematics, Princeton University, Princeton NJ 08544, USA

<sup>d</sup> Department of Mathematics, Rutgers University, Piscataway NJ 08854-8019, USA

## ARTICLE INFO

### Article history:

Received 31 December 2009

Available online xxx

Dedicated to Nihat Berker on the occasion of his 60th birthday

### Keywords:

Quenched disorder

Lattice spin systems

## ABSTRACT

The rounding of first-order phase transitions by quenched randomness is stated in a form which is applicable to both classical and quantum systems: The free energy, as well as the ground state energy, of a spin system on a  $d$ -dimensional lattice is continuously differentiable with respect to any parameter in the Hamiltonian to which some randomness has been added when  $d \leq 2$ . This implies absence of jumps in the associated order parameter, e.g., the magnetization in the case of a random magnetic field. A similar result applies in cases of continuous symmetry breaking for  $d \leq 4$ . Some questions concerning the behavior of related order parameters in such random systems are discussed.

© 2010 Elsevier B.V. All rights reserved.

## 1. Introduction

The effect of quenched randomness on the equilibrium and transport properties of macroscopic systems is a subject of great theoretical and practical interest which is close to Nihat's heart. He and his collaborators [1–3] made important contributions to the study of the changes brought about by such randomness in phase transitions occurring in the pure (nonrandom) system. These effects can be profound in low dimensions. Their understanding evolved in a somewhat interesting way. As was argued by Imry, Ma and others [1,4,5], even a very small amount of randomness can change the nature of the transition in dimensions  $d \leq 2$ . The validity of the reasoning involved was contentious as it conflicted with the prediction of a dimensional-reduction principle, which was arrived at through perturbative calculations carried to all orders [6]. Rigorous work by Imbrie [7] and Bricmont and Kupiainen [8] showed that, in the discrete case, the rounding phenomenon does not extend beyond two dimensions, as given by the Imry–Ma argument. Aizenman and Wehr [9,10] provided a rigorous proof of the rounding effect in low dimensions, which although limited to classical spin systems included a large variety of types of randomness not previously considered. Although some questions have arisen recently about whether the effect should be equally general for quantum systems [11], we have recently shown [12] that with some modification the same proof can be extended to quantum systems. Here we present these results in a unified way, and discuss related issues which appear in specific examples of interest.


To formulate the results in a general, but not the most general way, we consider a lattice system of generalized spin variables  $\underline{\sigma} = \{\sigma_x\}$ ,  $x \in \mathbb{Z}^d$ , where  $\sigma_x$  is an operator in a finite dimensional Hilbert space or, in the classical case, a function taking values in a finite set. The Hamiltonian is of the form

$$\mathcal{H}_\epsilon = \mathcal{K}(\underline{\sigma}; \underline{J}) - h \sum_x \kappa_x - \epsilon \sum_x \eta_x \kappa_x = \mathcal{H}_0 - \epsilon \sum_x \eta_x \kappa_x \quad (1)$$

\* Corresponding author. Tel.: +1 7324452390x2268.

E-mail address: [rafaelgr@physics.rutgers.edu](mailto:rafaelgr@physics.rutgers.edu) (R.L. Greenblatt).





**Fig. 1.** Phase diagrams of (a) the simple Ising model and (b) the Ising lattice gas in two dimensions; thick lines are first-order phase transitions.

**Theorem 2.** For the systems described above, with the random fields  $\vec{\eta}_x$  having a rotation-invariant distribution, the free energy is continuously differentiable in  $\vec{h}$  at  $\vec{h} = 0$  whenever  $\epsilon \neq 0, d \leq 4$ .

As before, the proof for classical systems is found in Ref. [10] and the proof for quantum systems in Ref. [12].

## 2. Higher order odd correlations

It can be proven rigorously for the models described by (2) with purely ferromagnetic interactions ( $J_{x-y} \geq 0$ ) and  $\lambda = 0$  that in the pure ( $\epsilon = 0$ ) system, not only the magnetization  $\langle \sigma_x \rangle$  but also all odd correlations, e.g.  $\langle \sigma_x^{(3)} \sigma_y^{(3)} \sigma_z^{(3)} \rangle, x \neq y \neq z$ , are discontinuous at  $h = 0$  for  $T < T_c$  [15]. The same is presumably true quite generally, particularly for the QRFIM with  $\lambda \neq 0$ ; a discontinuity in the magnetization brings with it also a discontinuity in all odd correlation functions. We may now ask whether the randomness in the external field, corresponding to  $\epsilon \neq 0$ , which removes the discontinuity in  $\langle \sigma_x \rangle$  also removes the discontinuity in all the odd correlation functions when averaged with respect to the randomness.

For the cases covered by (2) when  $J_{x-y} \geq 0$ , this question can be answered in the affirmative, including for  $\lambda \neq 0$ . In this case the system satisfies the Fortuin–Kasteleyn–Ginibre (FKG) inequalities [16]. Using these, it is possible to show that whenever all Gibbs states agree on the magnetization per site  $\langle \sigma_x \rangle$  (i.e. when the spontaneous magnetization  $M$  is zero) they also agree on all other local observables (the argument for random systems is spelled out in Ref. [17]). We do not have any such result for the case when  $J_{x-y}$  in (2) contains also some negative terms. We expect of course that this will be the case more generally, but we lack a proof.

For an instructive example let us consider the phase diagram of the Ising model (2) with  $\lambda = 0$  and  $d = 2$  with nearest neighbor interactions  $J_{\hat{e}} = J > 0$  ( $\hat{e} \in \{(0, 1), (1, 0)\}$ ),  $\epsilon = 0$  (see Fig. 1a). The free energy is analytic everywhere except on the bold part of the  $T$  axis. Along that line the free energy is continuous but its derivatives are discontinuous in all directions transverse to the  $T$  axis. That is, the free energy surface has a seam, i.e. a line of ‘kink singularities’.

Those singularities will be rounded through the addition of arbitrarily small randomness to the field  $h$ . However, the kink singularity would not be eliminated by small randomness in the bond strengths. That is, with  $J$  replaced by  $J + \epsilon b_{x,\hat{e}}$  with the  $b_{x,\hat{e}}$  i.i.d. random variables, there would still be spontaneous magnetization at low temperatures, though  $T_c$  may change, and so possibly will the nature of the singularity in  $\mathcal{F}$  at  $T_c$ . The randomness is tangential to the seam, corresponding to ‘orthogonal randomness’ in the terminology of Ref. [2].

Let us now consider this system in terms of lattice gas variables. Setting  $\xi_x = \frac{1}{2}(1 + \sigma_x) \in \{0, 1\}$ , the nonrandom Hamiltonian now takes the form

$$\mathcal{H}_0 = -4J \sum_{\langle x,y \rangle} \xi_x \xi_y - \mu \sum_x \xi_x + \text{constant}. \quad (8)$$

The chemical potential  $\mu$  is related to the parameters of the spin model by  $\mu = 2h - 8J$ , and so the phase diagram is now of the form shown in Fig. 1b. At the bold line the free energy has a discontinuous derivative in both variables, and randomness in either  $J$  or  $\mu$  will eliminate this first-order transition, rounding the seam in  $\mathcal{F}(T, \mu, 0)$ . To avoid destroying the first-order transition we have to add randomness to  $J$  and  $\mu$  in a correlated manner, so as to make it ‘tangential’ to the seam, i.e., in the form given in (2) in the  $\sigma$ -variables.

Similar reasoning can be applied quite generally. In particular if we consider the free energy  $\mathcal{F}(T, J, h, \epsilon = 0)$  for the system described by (2) with  $J$  containing interaction parameters multiplying both even and odd functions of the spins then, for  $T < T_c$ ,  $\mathcal{F}$  will have ‘multidimensional seams’ in directions transverse to all the odd terms – which is why they are all discontinuous in the absence of randomness. Adding then randomness to any odd term, e.g. to the one spin term as in (2), then rounds the discontinuity in all of them. If on the contrary we add noise only to the even terms, this is tangential to the ‘seams’ and doesn’t destroy the first-order transition.

This is also the case when adding noise in the QRFIM (2) to the transverse component of the field, i.e. letting  $\lambda \sum \sigma_x^{(1)} \rightarrow \sum (\lambda + \epsilon \eta_x) \sigma_x^{(1)}$ . As proven in Ref. [18] this does not destroy the discontinuity in  $\langle \sigma_x^{(3)} \rangle$ .

The situation is similar in the case when the spin at site  $x$ ,  $S_x$ , assumes the values  $S_x = -1, 0, 1$ . A particular version of such a system is the Blume–Capel model, whose nonrandom version is defined by

$$\mathcal{H}_0 = -J \sum_{\langle x,y \rangle} (S_x - S_y)^2 - \Delta \sum_x S_x^2 - h \sum_x S_x. \quad (9)$$




Fig. 2. Sections of the phase diagram for the Blume–Capel model.

The low temperature phase diagram of this system is known rigorously from Pirogov–Sinai theory [19,20]. A section is shown in Fig. 2 alongside the corresponding diagram at zero temperature.

In analogy with the Ising case, randomizing the magnetic field  $h$ , by using  $h \mapsto h + \epsilon \eta_x$ , will destroy all the first-order lines for  $d \leq 2$  (for  $d = 1$  this would be relevant only at  $T = 0$ ). Randomness in  $J$  or  $\Delta$  on the other hand should not destroy the discontinuity along the  $h = 0$  line. This is in accord with the analysis in Ref. [1]. Similar results will hold for the quantum version of this model with  $S_x$  now interpreted as  $S_x^{(3)}$ , and a transverse field term  $\sum_x S_x^{(1)}$  added.

### 3. Effects of bond randomness on the discontinuity of spontaneous magnetization

While the ‘tangential disorder’, in the sense discussed above, will not eliminate the discontinuity in the order parameter associated with the transverse field, e.g. the spontaneous magnetization at low enough temperatures, it *may* eliminate the possible discontinuity of the spontaneous magnetization if one was present at  $T_c$  in the non-random model. A general statement to that effect was proposed in Ref. [1].

An relevant example is provided by the two-dimensional Potts models with nearest neighbor ferromagnetic interactions. At  $Q > 4$  these models exhibit a discontinuity in the spontaneous magnetization at  $T_c$ , which coincides with a jump in the energy density. While Theorem 1 assures that randomization of the bond strengths will eliminate the energy discontinuity, it does not address the persistence or lack of it for the discontinuity in the spontaneous magnetization. In Ref. [1] it was argued that also the latter discontinuity will be eliminated, but this remains an interesting open challenge from the rigorous point of view [21].

### 4. One-dimensional, inverse square interaction

Another class of models exhibiting a jump in the spontaneous magnetization are the one-dimensional  $Q$ -state Potts models, at  $Q \geq 1$ , with ferromagnetic pair interactions  $J_{x-y}$  which asymptotically behave as  $J^+ / |x - y|^2$ . It is known that such systems have a phase transition, and that these exhibit somewhat unusual features [22,23] (and references therein). Among those is the discontinuity of the spontaneous ‘magnetization’  $M(T)$  at zero ‘magnetic’ field, which jumps at  $T_c$  from zero to a value satisfying  $M^2 J^+ / T_c \geq 1$ .

At the moment it is not clear to us whether for the  $1/r^2$  Potts models the discontinuity in  $M(T)$  may coincide with a discontinuity in the energy (it does not for  $Q = 2$  [24,25]). That will of course correspond to a jump at  $T_c$  in the derivative of  $\mathcal{F}$  with respect to  $T$ . If that occurs then the system will have at  $T = T_c$  at least  $Q + 1$  phases. In the absence of energy discontinuity one may expect that at the critical temperature the systems will have  $Q$  ordered phases in analogy with the results of Ref. [15] for Ising systems.

Let us consider now the effects of the additions of a random symmetry-breaking ‘magnetic’ field and of randomization, at suitably decaying rates, of the spin–spin interaction terms. As mentioned in the Introduction, the proof of the rounding phenomenon [10] applies also to the long range  $d = 1$  models as long as the pair interaction decays faster than  $|x - y|^{-3/2+\delta}$  (this ensures that the energy of interaction between an interval of size  $L$  and the rest of the system is less than  $L^{1/2}$ ). Thus, in the presence of a random magnetic field the spontaneous magnetization  $M(T)$  will vanish for all  $T$ . Randomization of the pair interaction would remove the discontinuity in the energy, if there was one, but unlike for the case discussed above we do not expect bond randomness to remove the discontinuity in  $M(T)$  at the (possibly altered) critical temperature. The reason for the possible difference with the two-dimensional models is the bound  $M^2 J^+ / T \geq 1$  which relates the discontinuity in the  $d = 1$  models to the purely long range part of the interaction.

### Acknowledgements

It has been both instructive and a personal pleasure for us to know Nihat Berker. We (MA and JLL) thank him for the many interesting discussions of physics as well as for his kind hospitality, which we enjoyed on different occasion at MIT and in Istanbul. We wish you Nihat many fruitful and enjoyable years past this birthday.

The first author's work was supported by NSF Grant DMR-044-2066 and AFOSR Grant AF-FA9550-04, the second author's work was supported by NSF Grant DMS-060-2360 and the third author's work was supported by NSF Grant DMR-044-2066 and AFOSR Grant AF-FA9550-04.

## References

- [1] K. Hui, A.N. Berker, Random-field mechanism in random-bond multicritical systems, *Phys. Rev. Lett.* 62 (21) (1989) 2507–2510.
- [2] A.N. Berker, Harris criterion for direct and orthogonal quenched randomness, *Phys. Rev. B* 42 (13) (1990) 8640–8642.
- [3] A. Malakis, A.N. Berker, I. Hadjiagapiou, N. Fytas, T. Papakonstantinou, Uncovering the secrets of the 2d random-bond Blume–Capel model, *Physica A*, in this issue (doi:10.1016/j.physa.2009.12.064).
- [4] Y. Imry, S.-k. Ma, Random-field instability of the ordered state of continuous symmetry, *Phys. Rev. Lett.* 35 (21) (1975) 1399–1401.
- [5] A. Aharony, Y. Imry, S.-k. Ma, Lowering of dimensionality in phase transitions with random fields, *Phys. Rev. Lett.* 37 (20) (1976) 1364–1367.
- [6] G. Parisi, N. Sourlas, Random magnetic fields, supersymmetry, and negative dimensions, *Phys. Rev. Lett.* 43 (11) (1979) 744–745.
- [7] J.Z. Imbrie, Lower critical dimension of the random-field Ising model, *Phys. Rev. Lett.* 53 (18) (1984) 1747–1750.
- [8] J. Bricmont, A. Kupiainen, Lower critical dimension for the random-field Ising model, *Phys. Rev. Lett.* 59 (16) (1987) 1829–1832.
- [9] M. Aizenman, J. Wehr, Rounding of first-order phase transitions in systems with quenched disorder, *Phys. Rev. Lett.* 62 (21) (1989) 2503–2506.
- [10] M. Aizenman, J. Wehr, Rounding effects of quenched randomness on first-order phase transitions, *Comm. Math. Phys.* 130 (3) (1990) 489–528.
- [11] P. Goswami, D. Schwab, S. Chakravarty, Rounding by disorder of first-order quantum phase transitions: Emergence of quantum critical points, *Phys. Rev. Lett.* 100 (1) (2008).
- [12] R.L. Greenblatt, M. Aizenman, J.L. Lebowitz, Rounding of first order transitions in low-dimensional quantum systems with quenched disorder, *Phys. Rev. Lett.* 103 (19) (2009) 197201.
- [13] T. Senthil, Properties of the random-field Ising model in a transverse magnetic field, *Phys. Rev. B* 57 (14) (1998) 8375–8380.
- [14] P. Vuilleumot, Thermodynamics of quenched random spin systems and applications to the problem of phase transitions in magnetic spin glass, *J. Phys. A: Math. Gen.* 10 (8) (1977) 1319.
- [15] J.L. Lebowitz, Coexistence of phases in Ising ferromagnets, *J. Stat. Phys.* 16 (6) (1977) 463–476.
- [16] C. Fortuin, P. Kasteleyn, J. Ginibre, Correlation inequalities on some partially ordered sets, *Comm. Math. Phys.* 22 (2) (1971) 89–103.
- [17] A. Bovier, *Statistical Mechanics of Disordered Systems: A Mathematical Perspective*, Cambridge University Press, 2006.
- [18] M. Campanino, A. Klein, J. Perez, Localization in the ground state of the Ising model with a random transverse field, *Comm. Math. Phys.* 135 (3) (1991) 499–515.
- [19] S. Pirogov, Y. Sinai, Phase diagrams of classical lattice systems, *Theoret. Math. Phys.* 25 (3) (1975) 1185–1192.
- [20] S. Pirogov, Y. Sinai, Phase diagrams of classical lattice systems continuation, *Theoret. Math. Phys.* 26 (1) (1976) 39–49.
- [21] L. Chayes, J.L. Lebowitz, V. Marinov, Percolation phenomena in low and high density systems, *J. Stat. Phys.* 129 (2007) 567–585.
- [22] M. Aizenman, J. Chayes, L. Chayes, C. Newman, Discontinuity of the magnetization in one-dimensional  $1/|x - y|^2$  Ising and Potts models, *J. Stat. Phys.* 50 (1) (1988) 1–40.
- [23] J. Imbrie, C. Newman, An intermediate phase with slow decay of correlations in one dimensional  $1/|x - y|^2$  percolation, Ising and Potts models, *Comm. Math. Phys.* 118 (2) (1988) 303–336.
- [24] P.W. Anderson, G. Yuval, Some numerical results on the Kondo problem and the inverse square one dimensional Ising model, *J. Phys. C* 4 (1971) 607–620.
- [25] J. Bhattacharjee, S. Chakravarty, J.L. Richardson, D.J. Scalapino, Some properties of a one-dimensional Ising chain with an inverse-square interaction, *Phys. Rev. B* 24 (1981) 3862–3865.