

On the number of lattice points between two enlarged and randomly shifted copies of an oval

Zheming Cheng¹, Joel L. Lebowitz², Péter Major³

¹Rutgers University, Department of Mathematics, New Brunswick, NJ 08903, USA

²Rutgers University, Department of Mathematics and Physics, New Brunswick, NJ 08903, USA

³Mathematical Institute of the Hungarian Academy of Sciences, P.O.B. 127, H-1364, Budapest, Hungary

Received: 29 October 1993/ 3 January 1994

Summary. Let A be an oval with a nice boundary in \mathbb{R}^2 , R a large positive number, $c > 0$ some fixed number and α a uniformly distributed random vector in the unit square $[0, 1]^2$. We are interested in the number of lattice points in the shifted annular region consisting of the difference of the sets $\{(R + c/R)A - \alpha\}$ and $\{(R - c/R)A - \alpha\}$. We prove that when R tends to infinity, the expectation and the variance of this random variable tend to $4c$ times the area of the set A , i.e. to the area of the domain where we are counting the number of lattice points. This is consistent with computer studies in the case of a circle or an ellipse which indicate that the distribution of this random variable tends to the Poisson law. We also make some comments about possible generalizations.

Mathematics Subject Classifications (1991): 60F5, 60K4, 11K06

1 Introduction

Using computer simulation Cheng and Lebowitz [4] studied the distribution of the number of lattice points in the domain between two concentric circles of radii $R + c/R$ and $R - c/R$ whose center is uniformly distributed on the unit square $[0, 1]^2$. (By lattice points we mean points from \mathbb{Z}^2 , i.e. from the set of points in \mathbb{R}^2 with integer coordinates.) This computer study, motivated by works of Sinai [9, 10], Bleher [2] and Major [8], suggested that for large R this distribution is asymptotically Poissonian with parameter $4\pi c$, i.e. with the area of the domain where we are counting the number of lattice points. A first step to check the correctness of this statement is to investigate whether the variance of this distribution is asymptotically $4\pi c$, i.e. whether the variance behaves as the Poissonian limit suggests. We answer this question in the affirmative. A similar statement holds for a class of ovals defined as follows.

Definition of an oval. A closed bounded convex set \mathbf{A} is an oval in \mathbb{R}^2 if it contains the origin in its interior, and its boundary is a smooth four times differentiable Jordan curve whose curvature is positive at all points.

We also introduce the following notations. Let $|\mathbf{A}|$ denote the area of a measurable set \mathbf{A} in \mathbb{R}^2 . Given some set $\mathbf{A} \subset \mathbb{R}^2$, $\alpha \in \mathbb{R}^2$ and number $R \in \mathbb{R}^+$ define the set $R\mathbf{A} - \alpha$ as

$$R\mathbf{A} - \alpha = \{\mathbf{u} \in \mathbb{R}^2; \mathbf{u} = R\mathbf{v} - \alpha, \mathbf{v} \in \mathbf{A}\}$$

and for $c > 0$, $R^2 > c$ introduce the difference set

$$\mathbb{O}_R(c, \alpha) = [(R + c/R)\mathbf{A} - \alpha] \setminus [R - c/R)\mathbf{A} - \alpha]. \quad (1.1)$$

Clearly, $|\mathbb{O}_R(c, \alpha)| = 4c|\mathbf{A}|$ for all α . The following Theorem is the main result of this paper.

Theorem. Let \mathbf{A} be an oval, $c > 0$ some fixed positive number and α a uniformly distributed random variable on $[0, 1]^2$. For $R > \sqrt{c}$ let $\xi_R = \xi_R(\alpha)$ denote the number of lattice points in the set $\mathbb{O}_R(c, \alpha)$ defined in (1.1). Then

$$E \xi_R = 4c|\mathbf{A}|, \quad (1.2)$$

$$\lim_{R \rightarrow \infty} \text{Var} \xi_R = 4c|\mathbf{A}|. \quad (1.3)$$

The investigation of the number of lattice points in a domain is a popular subject in number theory. See e.g. [6] for a recent treatment or [7]. This problem also has physical motivations, relating to the investigation of the statistics of eigenvalues in a quantum system with an integrable classical Hamiltonian. For example, if \mathbf{A} is a circle, the lattice points \mathbf{n} label energy levels $E(\mathbf{n}; \alpha) = |\mathbf{n} - \alpha|^2$ of the Laplacian $-(\nabla - \alpha)^2$ on the unit torus. These energies can be thought of as points on the real line and their statistics can be studied. This problem seems to be very hard. An easier problem is to consider not a fixed α but a random one distributed uniformly on the unit square $[0, 1]^2$, and this is what we have done. A widely accepted conjecture in the physics community is that the distribution of levels is, for typical systems in this class, locally Poissonian [1], i.e. the statistics of the energy levels in the interval $[E, E + L]$, L is fixed and E is uniformly distributed in an interval $[0, T]$ with $T \rightarrow \infty$, behave like Poisson distributed points with density π . In our context the conjecture is the following:

Let $P(n; R)$ be the probability that there are exactly n lattice points in $\mathbb{O}_R(c, \alpha)$. Then

$$\lim_{R \rightarrow \infty} P(n; R) = p(n) \quad \text{with } p(n) = [(4c|\mathbf{A}|)^n / n!] e^{-4c|\mathbf{A}|}. \quad (1.4)$$

Such a result was proved by Sinai [8] and Major [9] for the number of lattice points in scaled annuli domains bounded by *very* random curves. Here all the randomness comes from α , so the proof of (1.4), if indeed it is true, is far from trivial. Our theorem proves that the limit of the first and second moment of $P(n; R)$ has the right behavior when $R \rightarrow \infty$. In fact, our result also shows that

the covariance of pairs of distinct random variables $\eta_R^{(j)} = \xi_{R+jc} - |\mathbb{O}_R|$ vanishes as $R \rightarrow \infty$. This suggests that $\lim_{c \rightarrow \infty} \lim_{R \rightarrow \infty} (\xi_R - |\mathbb{O}_R|) / \sqrt{|\mathbb{O}_R|}$ should be a Gaussian random variable. This is consistent with taking the large parameter limit of the Poissonian distribution, but may be valid more generally.

2 Proof of the Theorem

The proof of relation (1.2) is simple. We can write $E \xi_R(\alpha)$ as the sum of the probabilities

$$E \xi_R(\alpha) = \sum_{\mathbf{m} \in \mathbb{Z}^2} P(\mathbf{m} \in \mathbb{O}_R(c, \alpha)) = \sum_{\mathbf{m} \in \mathbb{Z}^2} |\mathbb{O}_R \cap ([0, 1]^2 - \mathbf{m})| = |\mathbb{O}_R|.$$

Because of (1.2) formula (1.3) is equivalent to the relation

$$\lim_{R \rightarrow \infty} E \xi_R(\alpha) (\xi_R(\alpha) - 1) = |\mathbb{O}_R|^2. \quad (2.1)$$

We claim that

$$E \xi_R(\alpha) (\xi_R(\alpha) - 1) = \sum_{\mathbf{m} \in \mathbb{Z}^2 \setminus \{0\}} |\mathbb{O}_R \cap (\mathbb{O}_R - \mathbf{m})|. \quad (2.2)$$

Indeed,

$$\begin{aligned} E \xi_R(\alpha) (\xi_R(\alpha) - 1) &= \sum_{\mathbf{m} \in \mathbb{Z}^2} \sum_{\mathbf{m}_1 \in \mathbb{Z}^2 \setminus \{\mathbf{m}\}} P(\mathbf{m} \in \mathbb{O}_R(c, \alpha), \mathbf{m}_1 \in \mathbb{O}_R(c, \alpha)) \\ &= \sum_{\mathbf{m} \in \mathbb{Z}^2} \sum_{\mathbf{m}_1 \in \mathbb{Z}^2 \setminus \{\mathbf{m}\}} |[0, 1]^2 \cap (\mathbb{O}_R - \mathbf{m}) \cap (\mathbb{O}_R - \mathbf{m}_1)| \\ &= \sum_{\mathbf{m} \in \mathbb{Z}^2} \sum_{\mathbf{m}_1 \in \mathbb{Z}^2 \setminus \{0\}} |([0, 1]^2 + \mathbf{m}) \cap \mathbb{O}_R \cap (\mathbb{O}_R - \mathbf{m}_1)| \\ &= \sum_{\mathbf{m} \in \mathbb{Z}^2 \setminus \{0\}} |\mathbb{O}_R \cap (\mathbb{O}_R - \mathbf{m})|. \end{aligned}$$

Hence to prove the Theorem it is enough to prove relation (2.1) with the help of relation (2.2). This requires a good estimate on the area of $\mathbb{O}_R \cap (\mathbb{O}_R - \mathbf{m})$. First we introduce some notations.

Let us denote the boundary of the set $t\mathbf{A}$ for $t > 0$, by γ_t , and let $\gamma = \gamma_1$. For some $\mathbf{x} \in \gamma_t$ let $\varphi(\mathbf{x})$ denote the angle of the vector \mathbf{x} and $\psi(\mathbf{x})$ the angle of the normal of the curve γ_t at \mathbf{x} (pointing outside of the domain $t\mathbf{A}$) with the vector $\mathbf{e}_1 = (1, 0)$. Given some $\mathbf{z} \in \mathbb{R}^2 \setminus \{0\}$ let $d(\mathbf{z}, t)$ denote the diameter of the set $t\mathbf{A}$ in the direction \mathbf{z} , i.e.

$$d(\mathbf{z}, t) = \max\{|\mathbf{z}_1 - \mathbf{z}_2|; \mathbf{z}_1 \in \gamma_t, \mathbf{z}_2 \in \gamma_t, \mathbf{z}_1 - \mathbf{z}_2 = \lambda \mathbf{z}, \text{ with some } \lambda > 0\}.$$

Let $\mathbf{z}_{d,r}(t)$ and $\mathbf{z}_{d,l}(t)$ be the end points of this maximal vector, i.e. $\mathbf{z}_{d,r}(t) \in \gamma_t$, $\mathbf{z}_{d,l}(t) \in \gamma_t$, $\mathbf{z}_{d,r}(t) - \mathbf{z}_{d,l}(t) = \lambda \mathbf{z}$ with $\lambda > 0$ and $|\mathbf{z}_{d,r}(t) - \mathbf{z}_{d,l}(t)| = d(\mathbf{z}, t)$. For $\mathbf{z} \in \mathbb{R}^2$ let \mathbf{z}^\perp denote the vector \mathbf{z} rotated by $+\pi/2$, and define $K_t^+(\mathbf{z})$ and $K_t^-(\mathbf{z})$ as the half planes whose boundary is the line going through the points $\mathbf{z}_{d,r}(t)$ and

$\mathbf{z}_{d,t}(t)$ and which are in the direction \mathbf{z}^\perp and $-\mathbf{z}^\perp$ of this line, respectively. For $\mathbf{z} \in \mathbb{R}^2$ and $0 < |\mathbf{z}| \leq d(\mathbf{z}, t)$ define the (unique) points $\mathbf{z}_{t,r}^+, \mathbf{z}_{t,l}^+ \in K_t^+(\mathbf{z}) \cap \gamma_t$ and $\mathbf{z}_{t,r}^-, \mathbf{z}_{t,l}^- \in K_t^-(\mathbf{z}) \cap \gamma_t$ such that $\mathbf{z}_{t,r}^+ - \mathbf{z}_{t,l}^+ = \mathbf{z}$ and $\mathbf{z}_{t,r}^- - \mathbf{z}_{t,l}^- = \mathbf{z}$.

For $\mathbf{z}_1, \mathbf{z}_2 \in \gamma_R$ define the function

$$F_R(\mathbf{z}_1, \mathbf{z}_2) = \frac{4c^2 |\mathbf{z}_1| |\mathbf{z}_2| \cos(\varphi(\mathbf{z}_1) - \psi(\mathbf{z}_1)) \cos(\varphi(\mathbf{z}_2) - \psi(\mathbf{z}_2))}{R^4 |\sin(\psi(\mathbf{z}_1) - \psi(\mathbf{z}_2))|}$$

and for some $\mathbf{z} \in \mathbb{R}^2$, $0 < |\mathbf{z}| \leq d(\mathbf{z}, R)$ the functions

$$f_R^+(\mathbf{z}) = F_R(\mathbf{z}_{R,l}^+, \mathbf{z}_{R,r}^+), \quad f_R^-(\mathbf{z}) = F_R(\mathbf{z}_{R,l}^-, \mathbf{z}_{R,r}^-).$$

For $\mathbf{A} \subset \mathbb{Z}^2$ and $\mathbf{B} \subset \mathbb{Z}^2$ define their sum

$$\mathbf{A} + \mathbf{B} = \{\mathbf{x} + \mathbf{y}, \mathbf{x} \in \mathbf{A}, \mathbf{y} \in \mathbf{B}\}$$

and

$$\mathbf{A}^{(2)}(R) = R\mathbf{A} + (-R\mathbf{A}).$$

We claim that

$$\int_{\mathbf{A}^{(2)}(R)} [f_R^+(\mathbf{z}) + f_R^-(\mathbf{z})] d\mathbf{z} = 16c^2 |\mathbf{A}|^2. \quad (2.3)$$

Put

$$h_R(\mathbf{z}) = |\mathbb{O}_R \cap (\mathbb{O}_R - \mathbf{z})|, \quad \mathbf{z} \in \mathbb{R}^2. \quad (2.4)$$

We then also claim that

$$\lim_{R \rightarrow \infty} \left\{ \sum_{\mathbf{m} \in \mathbb{Z}^2 \setminus \{0\}} h_R(\mathbf{m}) - \int_{\mathbf{A}^{(2)}(R)} [f_R^+(\mathbf{z}) + f_R^-(\mathbf{z})] d\mathbf{z} \right\} = 0. \quad (2.5)$$

Relations (2.2), (2.3) and (2.5) together imply (2.1) hence also the Theorem. To prove (2.3) we introduce the maps

$$G^\pm: \mathbf{A}^{(2)}(R) \setminus \{0\} \mapsto \gamma_R \times \gamma_R, \\ G^+(\mathbf{z}) = (\mathbf{z}_{R,l}^+, \mathbf{z}_{R,r}^+), \quad G^-(\mathbf{z}) = (\mathbf{z}_{R,l}^-, \mathbf{z}_{R,r}^-).$$

Observe that the set $\text{Int } G^+(\mathbf{A}^{(2)}(R) \setminus \{0\}) \cap \text{Int } G^-(\mathbf{A}^{(2)}(R) \setminus \{0\})$ is empty, the maps G^\pm are diffeomorphisms on $\text{Int } \mathbf{A}^{(2)}(R) \setminus \{0\}$, and $G^+(\mathbf{A}^{(2)}(R) \setminus \{0\}) \cup G^-(\mathbf{A}^{(2)}(R) \setminus \{0\})$ is $\gamma_R \times \gamma_R \setminus \{(\mathbf{z}, \mathbf{z}), \mathbf{z} \in \gamma_r\}$.

The inverses of the maps $G^\pm(\mathbf{z})$ have Jacobians $|\sin(\psi(\mathbf{z}_{R,r}^\pm) - \psi(\mathbf{z}_{R,l}^\pm))|$. To see this we make the following observation: Let $[\mathbf{z}_1, \mathbf{z}_1 + d\mathbf{z}_1]$ and $[\mathbf{z}_2, \mathbf{z}_2 + d\mathbf{z}_2]$ be two small curves on γ_R starting from some points \mathbf{z}_1 and \mathbf{z}_2 respectively. Then the inverse of the map G^\pm maps the set $[\mathbf{z}_1, \mathbf{z}_1 + d\mathbf{z}_1] \times [\mathbf{z}_2, \mathbf{z}_2 + d\mathbf{z}_2]$ approximately to $\mathbf{z}_2 - \mathbf{z}_1 + \Delta(d\mathbf{z}_1, d\mathbf{z}_2)$, where $\Delta(d\mathbf{z}_1, d\mathbf{z}_2)$ is a parallelogram with one vertex at the origin, whose sides are the vectors $d\mathbf{z}_1$ and $d\mathbf{z}_2$. The area of this parallelogram is $|d\mathbf{z}_1| |d\mathbf{z}_2| |\sin(\psi(\mathbf{z}_1) - \psi(\mathbf{z}_2))|$. We can approximate the area of the image of the above domain by the inverse

map $(G^\pm)^{-1}$ with the area of this parallelogram. Since this approximation gives only an error of order $o(|dz_1||dz_2|)$ the Jacobian has the form we have stated.

The above relations imply that

$$\begin{aligned} & \int_{\mathbf{A}^{(2)}(R)} [f_R^+(z) + f_R^-(z)] dz \\ &= \int_{\gamma_R \times \gamma_R} \frac{4c^2}{R^4} |z_1| |z_2| \cos(\psi(z_1) - \varphi(z_1)) \cos(\psi(z_2) - \varphi(z_2)) dz_1 dz_2 \\ &= \left[\frac{2c}{R^2} \int_{\gamma_R} |z| \cos(\psi(z) - \varphi(z)) dz \right]^2 \\ &= \left[2c \int_{\gamma_1} |z| \cos(\psi(z) - \varphi(z)) dz \right]^2 = 16c^2 |\mathbf{A}|^2, \end{aligned}$$

hence relation (2.3) holds. To prove relation (2.5) we need some geometrical facts formulated in relations (2.9) and (2.10) and a lemma about the value of $h_R(\mathbf{z})$. They will be proved in the next Section.

Lemma 1 *There is some $\varepsilon > 0$ such that the function $h_R(\mathbf{z})$ defined in (2.4) satisfies the following estimates:*

- a) For $1 \leq |\mathbf{z}| < \varepsilon R$, $h_R(\mathbf{z}) < \text{const.}/R|\mathbf{z}|$.
b) For all $0 < \eta \leq \varepsilon$ and $\eta R \leq |\mathbf{z}|(1 - \eta)d(\mathbf{z}, R)$.

$$R^2 \{h_R(\mathbf{z}) - [f_R^+(z) + f_R^-(z)]\} \rightarrow 0 \quad \text{as } R \rightarrow \infty,$$

and the convergence is uniform in \mathbf{z} .

c) Let us fix some positive constant $B > 0$. Then the following inequalities hold:

- (c1) $h_R(\mathbf{z}) < \text{const.}/(R^{3/2} \sqrt{d(\mathbf{z}, R) - |\mathbf{z}|})$ if $(1 - \varepsilon)d(\mathbf{z}, R) \leq |\mathbf{z}| \leq d(\mathbf{z}, R) - B/R$.
(c2) $h_R(\mathbf{z}) < (\text{const.}/R)$ if $d(\mathbf{z}, R) - (B/R) \leq |\mathbf{z}| \leq d(\mathbf{z}, R) + B/R$.

d) $h_R(\mathbf{z}) = 0$ if $|\mathbf{z}| > d(\mathbf{z}, R) + B/R$, and B is larger than c times the diameter of the set \mathbf{A} .

To prove relation (2.5) let us introduce the sets

$$\begin{aligned} D_1^\varepsilon(R) &= \{\mathbf{z} \in \mathbb{R}^2, 0 < |\mathbf{z}| < \varepsilon R\}, \\ D_2^\varepsilon(R) &= (1 - \varepsilon)\mathbf{A}^{(2)}(R) \setminus D_1^\varepsilon(R), \\ D_3^\varepsilon(R) &= (1 - B/R^2)\mathbf{A}^{(2)}(R) \setminus (1 - \varepsilon)\mathbf{A}^{(2)}(R), \\ D_4^\varepsilon(R) &= (1 + B/R^2)\mathbf{A}^{(2)}(R) \setminus (1 - B/R^2)\mathbf{A}^{(2)}(R). \end{aligned}$$

Define the discrete measure μ on the positive half-line $[0, \infty]$,

$$\mu([0, x]) = \{\text{the number of lattice points in the set } x\mathbf{A}^{(2)}\},$$

where $\mathbf{A}^{(2)}$ denotes $\mathbf{A}^{(2)}(R)$ with $R = 1$. Define also the signed measure

$$v([0, x]) = \mu([0, x]) - x^2 \text{Area}(\mathbf{A}^{(2)}).$$

If \mathbf{A} is an oval, then $\mathbf{A}^{(2)}$ is again an oval [3]. (This means that the boundary of $\mathbf{A}^{(2)}$ is again strictly convex, smooth, and has positive curvature at all points.) Hence the results known for ovals can be applied to $\mathbf{A}^{(2)}$. In particular, we can state because of a result of Colin de Verdière [5] that

$$|v([0, x])| < \text{const.} x^{2/3} \quad \text{for all } x > 1. \quad (2.6)$$

Let us also remark that the normals of γ_R in the points $\mathbf{z}_{d,r}(R)$ and $\mathbf{z}_{d,l}(R)$ satisfy the relation

$$\psi(\mathbf{z}_{d,r}(R)) = \psi(\mathbf{z}_{d,l}(R)) + \pi \quad (2.7)$$

for all $\mathbf{z} \in \mathbb{R}^2 \setminus \{0\}$, i.e. the normals in the points $\mathbf{z}_{d,r}(R)$ and $\mathbf{z}_{d,l}(R)$ point in opposite directions. The half-line $\lambda \mathbf{z}$, $\lambda > 0$, intersects the boundary of $\mathbf{A}^{(2)}(R)$ at distance $d(\mathbf{z}, R)$ from the origin. Hence Part c1 of lemma 1 bounds the value of $h_R(\mathbf{z})$ for $\mathbf{z} \in D_4^{\varepsilon}(R)$ and Part c2 bounds the value of $h_R(\mathbf{z})$ for $\mathbf{z} \in D_4^{\varepsilon}(R)$. By Part a of Lemma 1

$$\sum_{\mathbf{m} \in \mathbb{Z}^2 \cap D_1^{\varepsilon}(R) \setminus \{0\}} h_R(\mathbf{m}) < \text{const.} \frac{1}{R} \sum_{\substack{\mathbf{m} \in \mathbb{Z}^2 \\ 0 < |\mathbf{m}| < \varepsilon R}} \frac{1}{|\mathbf{m}|} < \text{const.} \varepsilon.$$

Since $R^2 (f_R^+(R\mathbf{z}) + f_R^-(R\mathbf{z}))$ is uniformly continuous in the set $(1/R)D_2^{\varepsilon}(R)$, hence

$$\sum_{\mathbf{m} \in \mathbb{Z}^2 \cap D_2^{\varepsilon}(R)} h_R(\mathbf{m}) - \int_{D_2^{\varepsilon}(R)} [f_R^+(\mathbf{z}) + f_R^-(\mathbf{z})] d\mathbf{z} \rightarrow 0 \quad \text{as } R \rightarrow \infty$$

by Part b of Lemma 1. Put

$$H_R(t) = \sup_{z \in \partial \mathbf{A}^{(2)}} h_R(\mathbf{z}),$$

where $\partial \mathbf{A}^{(2)}$ denotes the boundary of $\mathbf{A}^{(2)}$. By using Part c1 of Lemma 1, integrating by parts and applying (2.6) we can write

$$\begin{aligned} \sum_{\mathbf{m} \in \mathbb{Z}^2 \cap D_3^{\varepsilon}(R)} h_R(\mathbf{m}) &\leq \int_{(1-\varepsilon)R}^{(R-B/R)} H_R(x) \mu(dx) \\ &< \frac{\text{const.}^{(R-B/R)}}{R^{3/2}} \int_{(1-\varepsilon)R}^x \frac{x}{\sqrt{R-x}} dx + \frac{\text{const.}^{(R-B/R)}}{R^{3/2}} \int_{(1-\varepsilon)R}^{(R-B/R)} \frac{v(dx)}{\sqrt{R-x}} \\ &< \text{const.} \sqrt{\varepsilon} + \frac{\text{const.}}{R^{3/2}} \left[\frac{v([0, x])}{\sqrt{R-x}} \right]_{(1-\varepsilon)R}^{(R-B/R)} \\ &\quad + \text{const.} \int_{(1-\varepsilon)R}^{(R-B/R)} \frac{v([0, x])}{R^{3/2}(R-x)^{3/2}} dx \\ &< \text{const.} \left(\sqrt{\varepsilon} + R^{-1/3} + R^{-5/6} \int_{(1-\varepsilon)R}^{(R-B/R)} \frac{1}{(R-x)^{3/2}} dx \right) \\ &< \text{const.} [R^{-1/3} + \sqrt{\varepsilon}]. \end{aligned}$$

Similarly, by Part c2 of Lemma 1

$$\begin{aligned} \sum_{\mathbf{m} \in \mathbb{Z}^2 \cap D_4^+(R)} h_R(\mathbf{m}) &\leq \text{const.} \int_{(R-B/R)}^{(R+B/R)} \frac{1}{R} \mu(dx) \\ &\leq \text{const.} \int_{(R-B/R)}^{(R+B/R)} \frac{x}{R} dx + \frac{\text{const.}}{R} v([R-B/R, R+B/R]) \\ &< \text{const.} [(1/R) + R^{-1/3}] < \text{const.} R^{-1/3}. \end{aligned}$$

The above relations together with Part d of Lemma 1 imply that

$$\sum_{\mathbf{m} \in \mathbb{Z}^2 \setminus \{0\}} h_R(\mathbf{m}) - \int_{D_2^+(R)} [f_R^+(\mathbf{z}) + f_R^-(\mathbf{z})] d\mathbf{z} = O(\sqrt{\varepsilon} + R^{-1/3}). \quad (2.8)$$

We claim that

$$I_1 = \int_{D_1^+(R)} [f_R^+(\mathbf{z}) + f_R^-(\mathbf{z})] d\mathbf{z} = O(\varepsilon) \quad (2.8')$$

and

$$I_2 = \int_{A^{(2)}(R) \setminus (1-\varepsilon)A^{(2)}(R)} [f_R^+(\mathbf{z}) + f_R^-(\mathbf{z})] d\mathbf{z} = O(\sqrt{\varepsilon}). \quad (2.8'')$$

Since $\varepsilon > 0$ can be chosen arbitrary small the above relations imply (2.5).

In Sect. 3 we shall prove the following statements. There is some $\varepsilon > 0$ such that if $|\mathbf{z}| = uR$ with some $0 < u < \varepsilon$, then the normals of γ_R satisfy the inequality

$$|\psi(\mathbf{z}_{R,r}^\pm) - \psi(\mathbf{z}_{R,l}^\pm)| > \text{const.} \cdot u, \quad (2.9)$$

and if $|\mathbf{z}| = (1-u)d(\mathbf{z}, R)$ with some $0 < u < \varepsilon$, then

$$|\psi(\mathbf{z}_{R,r}^\pm) - \psi(\mathbf{z}_{R,l}^\pm) - \pi| > \text{const.} \cdot \sqrt{u}. \quad (2.10)$$

In the first case we get that

$$|f_R^+(\mathbf{z})| + |f_R^-(\mathbf{z})| < \text{const.} \frac{1}{R^2 u} \quad (2.11)$$

and in the second case

$$|f_R^+(\mathbf{z})| + |f_R^-(\mathbf{z})| < \text{const.} \frac{1}{R^2 \sqrt{u}}. \quad (2.12)$$

Thus we get, by integrating in a polar coordinate system and applying the estimate (2.11), that

$$I_1 < \text{const.} \frac{1}{R^2} \int_0^\varepsilon R^2 u \frac{1}{u} du < \text{const.} \varepsilon.$$

Relation (2.12) implies that for $(1-\varepsilon)R \leq t \leq R$

$$\begin{aligned} \oint_{t\partial A^{(2)}} [f_R^+(\mathbf{z}) + f_R^-(\mathbf{z})] d\mathbf{z} &\leq \text{const.} \oint_{t\partial A^{(2)}} (1-t)^{-1/2} R^{-3/2} d\mathbf{z} \\ &\leq \text{const.} \frac{1}{\sqrt{(1-t)R}}. \end{aligned}$$

Integrating first on the curves γ_t , $(1-\varepsilon)R < t < R$, we get that

$$I_2 < \text{const.} \int_0^{\varepsilon R} R^{-1/2} u^{-1/2} du < \text{const.} \sqrt{\varepsilon}.$$

Hence we proved the Theorem with the help of Lemma 1 and formulas (2.9) and (2.10).

3 Proof of Lemma 1

We shall need a result about convex sets in the proof. To formulate it we introduce some notations. Let us fix some vector $\mathbf{z} \in \mathbb{R}^2$, $\mathbf{z} \neq 0$. Introduce the coordinate system whose coordinate axis x is in the direction $\mathbf{z}/|\mathbf{z}|$ and the coordinate axis y is its rotation with $+\pi/2$, in the direction $\mathbf{z}^\perp/|\mathbf{z}^\perp|$. In this new coordinate system let $(x_0(t), y_0(t))$ and $(x_1(t), y_1(t))$ be the points of tangency of the curve γ_t with the line parallel to the vector \mathbf{z} in the half-spaces $K_t^+(\mathbf{z})$ and $K_t^-(\mathbf{z})$, respectively. For $y_1(t) < u < y_0(t)$ the line $y = u$ intersects the curve γ_t in the points $x_{l,t}(u)$ and $x_{r,t}(u)$, $x_{l,t}(u) < x_{r,t}(u)$. We shall prove the following

Lemma 2 *There are some positive constants $A > 0$, $0 < B_1 < B_2$ depending only on the curve γ such that*

$$B_1 \sqrt{t(y_0(t) - u)} < x_{r,t}(u) - x_0(t) < B_2 \sqrt{t(y_0(t) - u)},$$

$$B_1 \sqrt{t(y_0(t) - u)} < x_0(t) - x_{l,t}(u) < B_2 \sqrt{t(y_0(t) - u)},$$

$$\frac{B_1 \sqrt{t}}{\sqrt{y_0(t) - u}} < -\frac{d}{du} x_{r,t}(u) < \frac{B_2 \sqrt{t}}{\sqrt{y_0(t) - u}},$$

$$\frac{B_1 \sqrt{t}}{\sqrt{y_0(t) - u}} < -\frac{d}{du} x_{l,t}(u) < \frac{B_2 \sqrt{t}}{\sqrt{y_0(t) - u}},$$

if $y_0(t) - tA < u < y_0(t)$, and

$$B_1 \sqrt{t(u - y_1(t))} < x_{r,t}(u) - x_1(t) < B_2 \sqrt{t(u - y_1(t))},$$

$$B_1 \sqrt{t(u - y_1(t))} < x_1(t) - x_{l,t}(u) < B_2 \sqrt{t(u - y_1(t))},$$

$$\frac{B_1 \sqrt{t}}{\sqrt{u - y_1(t)}} < \frac{d}{du} x_{r,t}(u) < \frac{B_2 \sqrt{t}}{\sqrt{u - y_1(t)}},$$

$$\frac{B_1 \sqrt{t}}{\sqrt{u - y_1(t)}} < -\frac{d}{du} x_{l,t}(u) < \frac{B_2 \sqrt{t}}{\sqrt{u - y_1(t)}},$$

if $y_1(t) < u < y_1(t) + tA$.

Proof of Lemma 2 Let us first restrict our attention to the case $t = 1$. It is more convenient to work with the inverse of the function $x_{r,1}(u)$. Let $(x, g(x))$ be

a small part of the curve γ in the neighborhood of the point $(x_0(1), y_0(1))$, and let $\rho(x)$ be the curvature of the curve γ in the point $(x, g(x))$. We can write in an interval $[x_0(1), x_0(1) + \eta]$, $\eta > 0$,

$$\rho(x) = -\frac{[1 + g'(x)^2]^{3/2}}{g''(x)}.$$

Put $z(x) = g'(x)$. Since $z(x_0(1)) = 0$, the last relation implies that

$$\int_0^{z(x)} \frac{dt}{(1+t^2)^{3/2}} = -\int_{x_0(1)}^x \frac{1}{\rho(t)} dt$$

or

$$\frac{z(x)}{\sqrt{1+z^2(x)}} = -P(x)$$

with $P(x) = \int_{x_0(1)}^x [1/\rho(t)] dt$. Since the curvature of γ is separated both from zero and infinity, there are some constants $K_2 > K_1 > 0$ such that

$$K_1(x - x_0(1)) < P(x) < K_2(x - x_0(1)).$$

Since $z(x) = -P(x)/\sqrt{1-P^2(x)}$ the last relation implies that

$$C_1(x - x_0(1)) < -g'(x) < C_2(x - x_0(1))$$

with some $C_2 > C_1 > 0$ in an interval $x \in [x_0(1), x_0(1) + \eta]$. Since $g(x_0(1)) = y_0(1)$ we get by integrating that

$$-\frac{1}{2}C_2(x - x_0(1))^2 < g(x) - y_0(1) < -\frac{1}{2}C_1(x - x_0(1))^2.$$

These formulas imply that $x_{r,1}(u)$, the inverse of $g(x)$, satisfies Lemma 1 for $t=1$ in an interval $[y_0(1) - A, y_0(1)]$. The remaining statements of Lemma 1 for $t=1$ can be proved in the same way. The case of general $t > 0$ follows from the case $t=1$ because of the homogeneity properties of γ_t . \square

Now we turn to the proof of relations (2.9) and (2.10). Here again we can restrict our attention to the case $R=1$. Let us recall that the curvature $\rho(\mathbf{x})$ of γ in a point $\mathbf{x} \in \gamma$ and the angle of the normal $\psi(\mathbf{x})$ in this point satisfy the relation

$$\frac{d\psi(\mathbf{x})}{ds(\mathbf{x})} = \rho(\mathbf{x}), \quad (3.1)$$

where $s(\mathbf{x})$ is the length of the arc $(\mathbf{x}_0, \mathbf{x})$ of γ with some fixed $\mathbf{x}_0 \in \gamma$.

Under the conditions imposed for formula (2.9) the length of the arc $(\mathbf{z}_{l,r}^+, \mathbf{z}_{l,r}^-)$ is greater than u . Since $\rho(\mathbf{x})$ is bounded from below by a positive constant, we get formula (2.9) by integrating (3.1).

The proof of (2.10) is similar. Here we can apply formula (2.7). Because of this formula it is enough to show that

$$\begin{aligned} |\psi(\mathbf{z}_{i,l}^\pm) - \psi(\mathbf{z}_{d,l}(1))| &> \text{const.} \sqrt{u}, \\ |\psi(\mathbf{z}_{i,r}^\pm) - \psi(\mathbf{z}_{d,r}(1))| &> \text{const.} \sqrt{u}, \end{aligned} \quad (3.2)$$

under the conditions imposed for (2.10).

Given two vectors \mathbf{z}_1 and \mathbf{z}_2 let $\angle(\mathbf{z}_1, \mathbf{z}_2)$ denote the angle between them. For $\mathbf{z} \in \gamma_1$ let $n(\mathbf{z})$ denote the normal vector to the curve γ_1 at \mathbf{z} . We make the following observation: For any $\mathbf{z} \in \mathbb{R}^2 \setminus \{0\}$ consider the end points $\mathbf{z}_{d,l} = \mathbf{z}_{d,l}(1)$ and $\mathbf{z}_{d,r} = \mathbf{z}_{d,r}(1)$ of the interval with maximal length in \mathbf{A} in the direction of \mathbf{z} . The normal of γ_1 in these points cannot be almost orthogonal to \mathbf{z} . More explicitly, there is some $\eta > 0$ such that

$$-\frac{1}{2}\pi + \eta < \angle(-\mathbf{z}, n(\mathbf{z}_{d,l})) < \frac{1}{2}\pi - \eta. \quad (3.3)$$

(This statement is equivalent to the following one: If \mathbf{z} is a boundary point of the oval $\mathbf{A}^{(2)}$, then the vector \mathbf{z} cannot be almost orthogonal to the normal of the boundary of $\mathbf{A}^{(2)}$ in this point. The equivalence of these two statements follows from the following argument. The vector $\mathbf{z}_{d,l} - \mathbf{z}_{d,r}$ is on the boundary of $\mathbf{A}^{(2)}$, and it is parallel to \mathbf{z} . The normal of $\mathbf{A}^{(2)}$ in this point is parallel to $n(\mathbf{z}_{d,l})$. The proof of the second statement is simpler.)

We claim that under the conditions imposed for formula (2.10) the distance of the parallel lines going through the points $\mathbf{z}_{i,l}^\pm$ and $\mathbf{z}_{i,r}^\pm$ and through the points $\mathbf{z}_{d,l}(1)$ and $\mathbf{z}_{d,r}(1)$ is bigger than $\text{const.} \sqrt{u}$. Put $m_l = m_l(\mathbf{z}) = 1/\cos \angle(\mathbf{z}, n(\mathbf{z}_{d,l}))$ and $m_r = m_r(\mathbf{z}) = 1/\cos \angle(\mathbf{z}, n(\mathbf{z}_{d,r}))$. Then $m_r = -m_l$ by (2.7), and by (3.3) there is some $\infty > K > 0$ such that $-K < m_r(\mathbf{z}) < K$ for all $\mathbf{z} \in \mathbb{R}^2 \setminus \{0\}$.

Let us fix a new coordinate system with the origin in a point of the line going through the points $\mathbf{z}_{i,r}$ and $\mathbf{z}_{d,r}$, with the x axis in the direction of the vector \mathbf{z} and y axis in the direction \mathbf{z}^\perp , and let us work in it. For $\varepsilon > v > 0$ let $\mathbf{z}_i^+(v) = (z_i^+(v), v)$ and $\mathbf{z}_r^+(v) = (z_r^+(v), v)$, $z_i^+(v) < z_r^+(v)$, be the two intersections of γ and the line $y = v$, and put $\mathbf{z}(v) = \mathbf{z}_i^+(v) - \mathbf{z}_r^+(v)$. It follows from Lemma 2 (with its application in the coordinate system with coordinate axes parallel to the normal and to the tangent vector of the curve γ_1 in the points $\mathbf{z}_{d,l}(1)$ and $\mathbf{z}_{d,r}(1)$, respectively) that there is some $C > 0$ such that $z_i^+(v) < z_{d,l}(1) + vm_l + Cv^2$ and $z_r^+(v) > z_{d,r}(1) + vm_l - Cv^2$. The above relations imply that $|\mathbf{z}(v)| = z_r^+(v) - z_i^+(v) > d(\mathbf{z}, 1) - 2Cv^2$. Since we imposed the condition $|\mathbf{z}| = d(\mathbf{z}, 1) - ud(\mathbf{z}, 1)$, this relation implies that the distance between the parallel lines going through the points $\mathbf{z}_{i,l}^\pm$ and $\mathbf{z}_{i,r}^\pm$ and through the points $\mathbf{z}_{d,l}(1)$ and $\mathbf{z}_{d,r}(1)$ is greater than $\text{const.} \sqrt{u}$. The same statement holds if $\mathbf{z}_{i,l}^\pm$ and $\mathbf{z}_{i,r}^\pm$ are replaced by $\mathbf{z}_{i,l}^-$ and $\mathbf{z}_{i,r}^-$. Hence the arcs $\mathbf{z}_{i,l}^\pm$, $\mathbf{z}_{d,l}$ and the arcs $\mathbf{z}_{i,r}^\pm$, $\mathbf{z}_{d,r}$ are longer than $\text{const.} \sqrt{u}$. Hence relation (3.1) and the strict positivity of $\rho(\mathbf{x})$ imply formula (3.2) and hence formula (2.10) too.

Now we turn to the proof of Lemma 1. We introduce the abbreviation R^+ instead of $(R + c/R)$ and R^- instead of $(R - c/R)$.

Proof of Lemma 1 Proof of Part a Let us introduce the coordinate system with x axis parallel to \mathbf{z} and y axis parallel to \mathbf{z}^\perp . We shall estimate, by means of Lemma 2, the length of the intersection of the set $\mathbb{O}_R \cap (\mathbb{O}_R - \mathbf{z})$ with the lines $y = u$ for different u 's.

Define $u_{R^+}(\mathbf{z})$ as the (unique) solution of the equation

$$x_{r,R^+}(u) - x_{l,R^+}(u) = |\mathbf{z}|, \quad u > u_{d,R^+},$$

where u_{d,R^\pm} is defined by the formula

$$x_{r,R^\pm}(u_{d,R^\pm}) - x_{l,R^\pm}(u_{d,R^\pm}) = \max_u [x_{r,R^\pm}(u) - x_{l,R^\pm}(u)],$$

that is, it is the level u at which the horizontal line $y = u$ has the longest intersection with the set $R^\pm \mathbf{A}$.

We claim that there is some $K > 0$ such that

$$x_{r,R^+}(u) - x_{l,R^+}(u) < |\mathbf{z}| + \text{const.}/|\mathbf{z}| \quad \text{if } [u_{R^+}(\mathbf{z}) - K/R] \leq u \leq u_{R^+}(\mathbf{z}), \quad (3.4)$$

$$x_{r,R^-}(u) - x_{l,R^-}(u) > |\mathbf{z}| \quad \text{if } u_{d,R^-} \leq u \leq u_{R^+}(\mathbf{z}) - K/R, \quad (3.4')$$

and

$$\left. \begin{array}{l} x_{r,R^+}(u) - x_{r,R^-}(u) < \frac{1}{2} \\ x_{l,R^+}(u) - x_{l,R^-}(u) < \frac{1}{2} \end{array} \right\} \quad \text{if } u_{d,R^-} \leq u \leq u_{R^+}(\mathbf{z}) - K/R. \quad (3.4'')$$

First we show that relations (3.4)–(3.4'') imply that

$$|\mathbb{O}_R \cap (\mathbb{O}_R - \mathbf{z}) \cap K_R^\pm(\mathbf{z})| < \text{const.}/(R|\mathbf{z}|). \quad (3.5)$$

To see this we show that the intersection of $\mathbb{O}_R \cap (\mathbb{O}_R - \mathbf{z})$ with the line $y = u$ has a length smaller than $\text{const.}/|\mathbf{z}|$ if $[u_{R^+}(\mathbf{z}) - K/R] \leq u \leq u_{R^+}(\mathbf{z})$, and it is empty if $u > u_{R^+}(\mathbf{z})$ or $u_{d,R^-} \leq u \leq u_{R^+}(\mathbf{z}) - K/R$. The above relations imply (3.5).

The above intersections are contained in the interval $[x_{l,R^+}(u), x_{r,R^+}(u) - |\mathbf{z}|]$ whose length is less than $\text{const.}/|\mathbf{z}|$ if $[u_{R^+}(\mathbf{z}) - K/R] \leq u \leq u_{R^+}(\mathbf{z})$ by (3.4). The distance $x_{r,R^+}(u) - x_{l,R^+}(u)$ is less than $|\mathbf{z}|$ for $u > u_{R^+}(\mathbf{z})$, because it is a (convex) monotone decreasing function of u in the interval $[u_{d,R^+}, y_0(R^+)]$ and it equals $|\mathbf{z}|$ for $u_{R^+}(\mathbf{z})$. This implies that the intersection $\mathbb{O}_R \cap (\mathbb{O}_R - \mathbf{z}) \cap \{(x, u), x \in \mathbb{R}^1\}$ is empty for $u > u_{R^+}(\mathbf{z})$. To see that it is empty for $u_{d,R^-} \leq u \leq u_{R^+}(\mathbf{z}) - K/R$ too, observe first that the intersection of \mathbb{O}_R with the line $y = u$ consists of two intervals, $[x_{l,R^+}(u), x_{l,R^-}(u)]$ and $[x_{r,R^-}(u), x_{r,R^+}(u)]$. The intersections

$$[x_{l,R^+}(u), x_{l,R^-}(u)] \cap ([x_{l,R^+}(u), x_{l,R^-}(u)] - |\mathbf{z}|)$$

and

$$[x_{r,R^-}(u), x_{l,R^+}(u)] \cap ([x_{r,R^-}(u), x_{l,R^+}(u)] - |\mathbf{z}|)$$

are empty because of (3.4'') and the condition $|\mathbf{z}| > 1$. The intersection

$$[x_{l,R^+}(u), x_{l,R^-}(u)] \cap ([x_{r,R^-}(u), x_{l,R^+}(u)] - |\mathbf{z}|)$$

is empty because of (3.4') and the intersection

$$[x_{r,R^-(u)}, x_{l,R^+(u)}] \cap ([x_{l,R^+(u)}, x_{l,R^-(u)}] - |\mathbf{z}|)$$

is always empty. In such a way we deduced (3.5) from (3.4)–(3.4'').

Let us recall the following notation. The point $(x_0(R^+), y_R^+)$ is the point of tangency of the curve γ_{R^+} with the line parallel to \mathbf{z} in the half-space $K_{R^+}(\mathbf{z})$. To prove relations (3.4)–(3.4'') let us first observe that because of the first two relations in Lemma 2

$$B_1 R(y_0(R^+)) - u_{R^+}(\mathbf{z}) < |\mathbf{z}|^2 < B_2 R(y_0(R^+)) - u_{R^+}(\mathbf{z}).$$

In particular, for $|\mathbf{z}| \geq 1$, $y_0(R^+) - u_{R^+}(\mathbf{z}) > \text{const.}/R$. Hence, by the third and fourth relations in Lemma 2

$$C_1 \frac{R}{|\mathbf{z}|} < -\frac{d}{du}(x_{r,R^+(u)} - x_{l,R^+(u)}) < C_2 \frac{R}{|\mathbf{z}|} \quad \text{for } u_{R^+}(\mathbf{z}) - \frac{K}{R} < u < u_{R^+}(\mathbf{z}) \quad (3.6)$$

with some $C_2 > C_1 > 0$. Since $x_{r,R^+(u_{R^+}(\mathbf{z}))} - x_{l,R^+(u_{R^+}(\mathbf{z}))} = |\mathbf{z}|$ we get relation (3.4) by integrating the right-hand side of (3.6) in the interval $[u, u_{R^+}(\mathbf{z})]$.

Because of the left-hand side of (3.6) we can choose for any $D > 0$ a number $C = C(D) > 0$ such that for $u_0 = u_{R^+}(\mathbf{z}) - (C/R)$ $x_{r,R^+(u_0)} - x_{l,R^+(u_0)} > |\mathbf{z}| + D/|\mathbf{z}|$. We rewrite this relation by turning from γ_{R^+} to γ_{R^-} . In this calculation we exploit that $|\mathbf{z}| < \text{const. } R$ if $\mathbf{z} \in \gamma_{R^+}$. Putting $u_1 = (1 - \eta)u_0 = (1 - c/R^2)(1 + c/R^2)^{-1} u_0$ we have $u_1 > u_{R^+}(\mathbf{z}) - K/R$ with an appropriate $K > 0$, $\eta = O(R^{-2})$ and

$$\begin{aligned} x_{r,R^-(u_1)} - x_{l,R^-(u_1)} &= (1 - \eta)(x_{r,R^+(u_0)} - x_{l,R^+(u_0)}) \\ &> (1 - \eta)(|\mathbf{z}| + D/|\mathbf{z}|) > |\mathbf{z}| \end{aligned}$$

if $D > 0$ is sufficiently large. This means that relation (3.4') holds for $u_1 > u_{R^+}(\mathbf{z}) - K/R$. Because of the monotonicity of $x_{r,R^-(u_0)} - x_{l,R^-(u_0)}$ for $u_{d,R^-} < u < y_0(R^-)$ relation (3.4') holds.

To prove relation (3.4'') first we observe that $x_{r,R^+}(\bar{u}) = (1 + \beta)x_{r,R^-}(u)$ with $1 + \beta = (1 + c/R^2)(1 - c/R^2)^{-1}$ and $\bar{u} = (1 + \beta)u$. Hence $x_{r,R^+}(\bar{u}) = x_{r,R^-}(u) + O(1/R)$ and $|u - \bar{u}| < L/R$ with some $L > 0$. The derivative $\frac{d}{du} x_{r,R^+}(u)$ is a monotone decreasing function of u , hence it follows from the third relation in Lemma 2 that

$$\left| \frac{d}{du} x_{r,R^+}(u) \right| < \frac{R}{3L}$$

if $[y_1(R^+) - K/R] \leq u \leq [y_0(R^+) - K/R]$, and $K > 0$ is chosen sufficiently large. Hence

$$|x_{r,R^+}(u) - x_{r,R^+}(\bar{u})| < (R/3L)|u - \bar{u}| + O(1/R) < 1/2.$$

The first relation of (3.4'') is proved, and the second one can be proved in the same way.

We have proved relation (3.5). It can be proved in the same way if $K_{R^+}(\mathbf{z})$ is replaced by $K_{R^-}(\mathbf{z})$. These two relations together imply Part a.

Proof of Part b. We define two parallelograms $P^+(\mathbf{z})$ and $P^-(\mathbf{z})$. The parallelogram $P^+(\mathbf{z})$ is bounded by two pairs of parallel lines, the lines of the first pair are going through the points $(1+c/R^2)\mathbf{z}_{R,l}^+$ and $(1-c/R^2)\mathbf{z}_{R,l}^+$ and they have normal $\psi(\mathbf{z}_{R,l}^+)$, the lines of the second pair are going through the points $(1-c/R^2)\mathbf{z}_{R,r}^+-\mathbf{z}$ and $(1+c/R^2)\mathbf{z}_{R,r}^+-\mathbf{z}$ and they have normal $\psi(\mathbf{z}_{R,r}^+)$. The parallel pairs of lines bounding $P^-(\mathbf{z})$ are going through the points $(1-c/R^2)\mathbf{z}_{R,l}^-$ and $(1+c/R^2)\mathbf{z}_{R,l}^-$ with normal $\mathbf{z}_{R,l}^-(\psi)$, and through the points $(1-c/R^2)\mathbf{z}_{R,r}^- - \mathbf{z}$ and $(1+c/R^2)\mathbf{z}_{R,r}^- - \mathbf{z}$ with normal $\mathbf{z}_{R,r}^-(\psi)$. The parallelograms $P^+(\mathbf{z})$ and $P^-(\mathbf{z})$ have area $f_R^+(\mathbf{z})$ and $f_R^-(\mathbf{z})$, respectively, and they are disjoint if $|\mathbf{z}| > \eta R$. Since the difference of these parallelograms and the domains $\mathbb{O}_R \cap (\mathbb{O}_R - \mathbf{z}) \cap K_R^+(\mathbf{z})$ and $\mathbb{O}_R \cap (\mathbb{O}_R - \mathbf{z}) \cap K_R^-(\mathbf{z})$ have an area of order $o(R^{-2})$, the above relations imply Part b of Lemma 1.

Proof of Part c Let us work in the coordinate system with origin $z_{d,l}(R^+)$, with x -coordinate axis in the direction $-n(z_{d,l}(R^+))$, the normal of γ_{R^+} in the point $z_{d,l}(R^+)$ showing inside the domain \mathbf{A}_{R^+} , and with y -axis in the direction $\mathbf{z}_e = -n(z_{d,l}(R^+))^\perp$, the tangent of γ_{R^+} in this point which is obtained when the x axis is rotated with angle $+\pi/2$. Let $y_{R^+}^\pm(u)$ be the y coordinate of the intersection of the set $\gamma_{R^+} \cap K_{R^+}^\pm(\mathbf{z})$ with the line $x = u$ and $y_{R^+}^\pm(u)$ the y coordinate of the intersection of the set $\gamma_{R^+} \cap K_{R^+}^\pm(\mathbf{z})$ with this line. We shall estimate the length of the intersection of \mathbb{O}_R with the line $x = u$. We shall prove that the Lebesgue measure of this intersection satisfies the inequality

$$\lambda(\mathbb{O}_R \cap \{(u, y), y \in \mathbb{R}^{(1)}\}) < \text{const.} \max\left(\frac{1}{\sqrt{Ru}}, 1\right) \quad \text{if } 0 < u < \eta R \quad (3.7)$$

with some $\eta > 0$.

By Lemma 2

$$|y_{R^+}^\pm(u)| < \text{const.} \sqrt{Ru} \quad \text{if } 0 < u < AR,$$

and this inequality implies (3.7) in the case $u < K/R$ with some $K > 0$. We shall show, using again Lemma 2, arguing similarly as in the proof of relation (3.4) in the proof of Part a that

$$|y_{R^+}^\pm(u) - y_{R^+}^\mp(u)| < \text{const.} / \sqrt{Ru} \quad \text{if } (K/R) < u < \eta R, \quad (3.8)$$

which relation completes the proof of (3.7). To prove (3.8) we have to express

$$y_{R^+}^\pm(\cdot) \text{ by } y_{R^+}^\mp(\cdot) \text{ and to exploit that by Lemma 2 } \left| \frac{d}{dv} y_{R^+}^\pm(v) \right| < \text{const.} \sqrt{R}/\sqrt{u} \text{ if}$$

$K/R < u < v < u + K/R$.

Let $\mathbf{v}_{R^+} = (v_{R^+,1}^+, v_{R^+,2}^+)$ and $\mathbf{v}_{R^+}^- = (v_{R^+,1}^-, v_{R^+,2}^-)$ be the points of intersection of γ_{R^+} and $\gamma_{R^+} - \mathbf{z}$ in the half planes $K_{R^+}^+(\mathbf{z})$ and $K_{R^+}^-(\mathbf{z})$, respectively. We shall prove that there is some constant $K > 0$ such that

$$\mathbb{O}_R \cap (\mathbb{O}_R - \mathbf{z}) \cap K_{R^+}^+(\mathbf{z}) \cap \{(v, y), y \in \mathbb{R}^1\} = \emptyset \quad \text{if } v \notin \left[v_{R^+,1}^+ - \frac{K}{R}, v_{R^+,1}^+ + \frac{K}{R} \right] \quad (3.9)$$

and

$$\mathbb{O}_R \cap (\mathbb{O}_R - \mathbf{z}) \cap K_{R^+}(\mathbf{z}) \cap \{(v, y), y \in \mathbb{R}^1\} = \emptyset \quad \text{if } v \notin \left[v_{R^+,1} - \frac{K}{R}, v_{R^+,1} + \frac{K}{R} \right] \quad (3.9')$$

We also claim that under the conditions of Part c1 or Part c2 of Lemma 1

$$v_{R^+,1}^\pm > \text{const.}(d(\mathbf{z}, R) - |\mathbf{z}|). \quad (3.10)$$

Relations (3.7), (3.9), (3.9') and (3.10) together imply Part c of Lemma 1. To prove relation (3.10) let us consider the projection of the vectors $\mathbf{v}_{R^+}^\pm$ and $(d(\mathbf{z}, R) - |\mathbf{z}|)(\mathbf{z}/|\mathbf{z}|) - \mathbf{v}_{R^+}^\pm$ to the direction of the vector $-n(\mathbf{z}_{d,l}(R^+))$. The sum of these two vectors, which is the projection of $(d(\mathbf{z}, R) - |\mathbf{z}|)(\mathbf{z}/|\mathbf{z}|)$ to $-n(\mathbf{z}_{d,l}(R^+))$ is longer than $\text{const.}(d(\mathbf{z}, R) - |\mathbf{z}|)$ because of relation (3.3). On the other hand, the proportion of the length of these two vectors is separated both from zero and infinity because of relation (3.3) which implies this relation if the projection is done in the orthogonal direction $n(\mathbf{z}_{d,l}(R^+))^\perp$ and Lemma 2.

To prove relations (3.9) and (3.9') we introduce the following notation. Let $s_{R^+}^\pm(u)$ be the y coordinate of the intersection of the set $(\gamma_{R^+} - \mathbf{z}) \cap K_{R^+}^\pm(\mathbf{z})$ with the line $x = u$. Since $s_{R^+}^\pm(v_{R^+,2}^\pm) = y_{R^+}^\pm(v_{R^+,2}^\pm)$ we get by expression $s_{R^+}^\pm(\cdot)$ through $s_{R^+}^\pm(\cdot)$, exploiting the lower bound on the derivative of the function $s_{R^+}^\pm(\cdot)$ given by Lemma 2 and arguing similarly to the proof of relation (3.7) that $s_{R^+}^\pm(v) > x_{R^+}^\pm(v)$ or $v < 0$ if $v < v_{R^+,1}^\pm - K/R$ with some sufficiently large $K > 0$. If $v < 0$, then the set $\mathbb{O}_R \cap \{(v, y), y \in \mathbb{R}^1\}$ is empty. Hence

$$\mathbb{O}_R \cap \{(v, y), y \in \mathbb{R}^1\} = \emptyset, \quad \text{if } v < v_{R^+,1}^\pm - \frac{K}{R}.$$

By changing the role of γ_R and $\gamma_R - \mathbf{z}$ we get that

$$\mathbb{O}_R \cap \{(v, y), y \in \mathbb{R}^1\} = \emptyset, \quad \text{if } v > v_{R^+,1}^\pm + \frac{K}{R}.$$

The last two relations together imply (3.9). The proof of (3.9') is similar.

In such a way we have proved Part c of Lemma 1. The proof of Part d is trivial, since in this case even the set $R^+ \mathbf{A} \cap (R^+ \mathbf{A} - \mathbf{z})$ is empty. \square

4 Some concluding remarks

In this Section we discuss the conjecture about the Poissonian distribution of a randomly placed circle suggested by the computer study of Cheng and Lebowitz [4] and briefly explain what kind of approach is suggested by the present paper.

It is relatively easy to show that the Poissonian limit for the number of lattice points in $\mathbb{O}_R(c, \alpha)$ would follow from the following generalization of formula (2.1):

$$\lim_{R \rightarrow \infty} E \xi_R(\alpha) (\xi_R(\alpha) - 1) \cdots (\xi_R(\alpha) - k + 1) = |\mathbb{O}_R|^k = [4c|\mathbf{A}|]^k \quad \text{for all } k \geq 1. \quad (4.1)$$

Some modification of the argument in the proof of formula (2.2) gives that

$$E \xi_R(\alpha)(\xi_R(\alpha)-1)\cdots(\xi_R(\alpha)-k+1) \\ = \sum_{\substack{\mathbf{m}_1, \dots, \mathbf{m}_{k-1} \text{ are different} \\ \text{points in } \mathbb{Z}^2 \setminus \{0\}}} |\mathbb{O}_R \cap (\mathbb{O}_R - \mathbf{m}_1) \cap \cdots \cap (\mathbb{O}_R - \mathbf{m}_{k-1})|. \quad (4.2)$$

It is relatively simple to prove the identity

$$\int |\mathbb{O}_R \cap (\mathbb{O}_R - \mathbf{z}_1) \cap \cdots \cap (\mathbb{O}_R - \mathbf{z}_{k-1})| dz_1 \dots dz_{k-1} = [4c|\mathbf{A}|]^k. \quad (4.3)$$

Hence to prove the Poissonian limit it would be enough to show that for large R the replacement of the sum in (4.2) by the integral in (4.3) causes a negligible error for all $k=1, 2, \dots$. Actually, we proved this for $k=1$ and 2. But the proof for $k \geq 3$ is much harder. In our proof we exploited the independence caused by the random shift $\alpha \in [0, 1]^2$. But this independence is not sufficient for the proof of (4.1) if $k \geq 3$. In this case some deep number theoretical statement would be needed which states that certain functions of the k -tuples of lattice points $(\mathbf{m}_1, \dots, \mathbf{m}_k)$ are almost uniformly distributed. We could give an explicit formulation of this statement, but since this requires complicated notations and would lead to a problem that we cannot handle we omit it.

Finally, we only briefly mention that the higher dimensional version of the problem can be discussed similarly, and the natural modification of the result holds. Moreover, the same technique gives a stronger result, namely in the case of d -dimensional convex annular set the first d moments converge to the moments of a Poissonian random variable. This strengthening is related to the fact that the shift with a random vector which is uniformly distributed in the d -dimensional unit cube provides more randomness for large d .

Acknowledgement. The authors would like to thank E. Makai Jr. for some useful explanations on convex geometry. J.L. Lebowitz and P. Major would like to thank the IHES in Bures-Sur-Yvette where this work was completed. The work of Zh. Cheng and J.L. Lebowitz was supported in part by the NSF Grant DMR 89 18902.

References

1. Berry, M.V., Tabor, M.: Level clustering in the regular spectrum. Proc. R. Soc. Lond., Ser. A **356**, 375–394 (1977)
2. Bleher, P.M.: Quasi-classical expansion and the theory of quantum chaos. (Lect. Notes Math., vol. 1469, pp. 60–89) Berlin Heidelberg New York: Springer, 1991
3. Bonnesen, T., Fenchel, W.: Theorie der konvexen Körper. Ergebnisse der Mathematik 3. Bd. Heft 1, New York: Chelsea, 1948
4. Cheng, Zh., Lebowitz, L.J.: Statistics of energy levels in integrable quantum systems. Phys. Rev. A **44** (6), 3399–3402 (1991)
5. Colin de Verdière, Y.: Nombre de points entiers dans une famille homothétique de domaines de \mathbf{R}^n . Ann. Scien., Éc. Norm. Sup Sér., IV **10**, 559–576 (1977)
6. Huxley, M.N.: Exponential sums and lattice points. Proc. Lond. Math. Soc. **60** (3), 470–502 (1990).

7. Kendall, D.G.: On the number of lattice points inside a random oval. *Quart. J. Math. Oxford* **19** 1–26 (1948)
8. Major, P.: Poisson law for the number of lattice points in a random strip with finite area *Probab. Theory Relat. Fields* **92**, 423–464 (1992)
9. Sinai, Ya. G.: Poisson distribution in a geometric problem. In: Ya. G. Sinai (ed.) *Advances in Soviet Mathematics*, Vol. 3 199–214 AMS, Providence, 1991
10. Sinai, Ya. G.: *Mathematical problems in the theory of quantum chaos*. (Lect. Notes Math. vol. 1469, pp. 41–52) Berlin Heidelberg New York: Springer, 1991