

# Statistical Mechanics and Typical Behavior of Macroscopic Systems

Joel L. Lebowitz

Departments of Mathematics and Physics  
Rutgers University

## Abstract

Statistical mechanics predicts, with great certainty, behavior of individual macroscopic systems, both in equilibrium and out of it. This behavior is **typical** for systems represented by the usual microcanonical ensemble or those derived from it by constraining the microstates to regions corresponding to nonequilibrium macrostates. These take probability to be proportional to phase space volume in classical systems and to “number of wave functions” in quantum systems. I will illustrate this by describing Lanford’s derivation of the Boltzmann equation and by some recent results for the time evolution of an ideal gas.

---

I thank S. Goldstein, E. Speer, and O. Penrose for many helpful comments and A. Krueger for technical help with these slides. I also thank the NSF and the AFOSR for long term support.

# Classical Systems

In classical mechanics, the microstate of a system of  $N$  particles confined to a box  $V$  in  $\mathbb{R}^d$  is a point  $X$  in the  $2dN$ -dimensional phase space,  $\Gamma$ ,

$$X = (\mathbf{r}_1, \mathbf{v}_1, \dots, \mathbf{r}_N, \mathbf{v}_N), \quad \mathbf{r}_i \in V \subset \mathbb{R}^d, \quad \mathbf{v}_i \in \mathbb{R}^d \quad (1)$$

Its time evolution is given by a Hamiltonian  $H(X)$  which conserves energy, so  $X(t) = T_t X$  will be confined to the energy surface  $H(X) = E$ .

This time evolution,  $T_t X = X(t)$ , is reversible: given  $X = (\mathbf{r}_1, \mathbf{v}_1, \dots, \mathbf{r}_N, \mathbf{v}_N)$  define  $RX = (\mathbf{r}_1, -\mathbf{v}_1, \dots, \mathbf{r}_N, -\mathbf{v}_N)$  then

$$T_s R T_s X = RX. \quad (2)$$

A macroscopic system is one with “very large”  $N$ , say  $N \gtrsim 10^{20}$ . For such systems it is more appropriate to replace the energy surface by a thin shell surrounding that surface, to which I will always refer as  $\Gamma_E$ . The micro-canonical ensemble is a uniform measure on  $\Gamma_E$ .

For macroscopic systems an overwhelming majority of the microstates in the micro-canonical ensemble give the same “thermodynamic” behavior: such behavior is typical, i.e. it has probability close to one for this probability distribution.

In fact, the fraction of systems with noticeable macroscopic deviations from the average behavior, computed in such an ensemble, is exponentially small in the number of degrees of freedom of the system.

The functions on the phase space which correspond to such “thermodynamic” behavior are sums of terms which depend only on the coordinates and momenta of a few particles.

Once one accepts the applicability of these measures (ensembles) to physical systems they can be used to predict the typical behavior of individual macroscopic systems *and* not just some average behavior.

The observed behavior does not require explanations based on ergodicity, time averaging, or subjective information theory.

# Macrostates

To describe the macroscopic state of such a system of  $N$  particles in a box  $V$ , we make use of a much cruder description than that provided by the microstate  $X$ . We shall denote by  $M$  such a macroscopic description.

In general we can describe a macrostate  $M$  by specifying a set of macrovariables  $M(X) = \{M_1(X), M_2(X), \dots, M_n(X)\}$ , with resolution  $\Delta M = \{\Delta M_j\}$ . We identify these macrostates with the elements of a partition of the phase space into sets  $\Gamma$  of the form

$$\Gamma_M = \{X | M_j \leq M_j(X) \leq M_j + \Delta M_j, j = 1, \dots, n\}. \quad (3)$$

These provide a coarse-grained description in the sense that many different  $X$  correspond to the same  $\Gamma_M$ .

It can be shown<sup>1</sup> that for all reasonable choices of the  $M$ 's there is in every  $\Gamma_E$  of a macroscopic system one dominant region  $\Gamma_M$  which has most of the volume of  $\Gamma_E$ . This is called the equilibrium macrostate  $M_{\text{eq}}$ ,

$$\frac{|\Gamma_{M_{\text{eq}}}|}{|\Gamma_E|} = 1 - \varepsilon \quad (4)$$

with  $\varepsilon \ll 1$ .

A system in a microstate  $X$  is then in macroscopic thermal equilibrium relative to the choice of the  $M$ 's iff  $X \in \Gamma_{M_{\text{eq}}}$ . When  $M$  specifies a nonequilibrium state,  $|\Gamma_M|$  is very much smaller, see Figure 1 for a generic choice of reasonable  $M$ 's. This is essentially a consequence of the law of large numbers.

---

<sup>1</sup>La73




Figure 1: Schematic Picture of the decomposition of  $\Gamma_E$ . Here  $\Gamma_{\text{eq}} \equiv \Gamma_{M_{\text{eq}}}$ .

The second picture is slightly more faithful.

Neither shows the topology or differences in relative sizes of the different  $\Gamma_M$ 's. In general, the closer  $M$  is to  $M_{\text{eq}}$  the larger  $\Gamma_M$ .

All thermodynamic functions, not just those used to define the  $M$ 's, have approximately the same value for almost all  $X \in \Gamma_{M_{\text{eq}}}$ , hence these values are typical for the uniform measure on  $\Gamma_{M_{\text{eq}}}$  and observed for (almost) all systems in equilibrium.

In fact since  $|\Gamma_{M_{\text{eq}}}| \simeq |\Gamma_E|$  they are also typical of the microcanonical measure on  $\Gamma_E$ .

This justifies the use of the microcanonical ensemble for  $N \gg 1$  to compute relevant properties of an equilibrium system, independent of whether or not the dynamics is ergodic in a mathematical sense.

## Nonequilibrium States

Thus, most microstates in  $\Gamma_E$  of a macroscopic system correspond to the system being in equilibrium.

Fortunately there are also microstates which correspond to macroscopic systems which are out of equilibrium (or we would not be here).

Consider now the time evolution of the macro variables  $M(X(t))$  when the system is in the microstate  $X(t_0) \in \Gamma_M$ ,  $M \neq M_{\text{eq}}$ , at an “initial” time  $t_0$ , and is isolated for  $t > t_0$ .

## Approach to Equilibrium

Boltzmann (also Maxwell, Kelvin, ...) argued that given the disparity in the sizes of the  $\Gamma_M$  corresponding to the various macrostates, the evolution of a microstate  $X(t_0)$ , typical wrt a uniform measure on  $\Gamma_M$ , will be such that  $|\Gamma_{M(X(t))}|$  will not decrease (on a macroscopic scale) for  $t > t_0$ .

Thus the evolution towards equilibrium of macroscopic systems which start in the region  $\Gamma_M$ ,  $M \neq M_{\text{eq}}$ , and are kept (effectively) isolated afterwards, is typical with respect to the micro-canonical measure restricted to  $\Gamma_M$ .

N.B. We have assumed that such measures give the right probabilities for physical systems.

## Boltzmann's Entropy

To make a connection with the Second Law, Boltzmann defined the (Boltzmann) entropy of a macroscopic system in a microstate  $X$  as

$$S_B(X) = \log |\Gamma_{M(X)}| = S_B(M(X)). \quad (5)$$

The entropy of the equilibrium macrostate

$$S_B(M_{\text{eq}}) = \log |\Gamma_{M_{\text{eq}}}| \simeq \log |\Gamma_E| \quad (6)$$

Boltzmann then showed that  $S_B(M_{\text{eq}})$  agrees with the Clausius thermodynamic entropy of a gas in equilibrium.

Boltzmann's heuristic argument for the non-decrease of entropy, based on relative phase space volume, is the correct explanation for the behavior typically observed in actual macroscopic systems. It is, however, very far from a mathematical theorem and contains no quantitative information about time scales<sup>2</sup>.

A proof would be provided by the rigorous derivation from the microscopic dynamics of the kinetic and hydrodynamic equations<sup>3</sup> commonly used to describe the time asymmetric, entropy increasing behavior of macroscopic systems out of equilibrium.

This has been achieved so far only for the Boltzmann equation for dilute gases. This was done rigorously (in appropriate limits) by Oscar Lanford in 1975.

---

<sup>2</sup>As Mark Kac said: an argument convinces a reasonable person, a proof convinces a stubborn one.

<sup>3</sup>such as the heat equation, Navier-Stokes equations, etc.

# The Boltzmann Equation for Dilute Gases

Following Boltzmann, we refine the description of a macrostate  $M$  by noting that the microstate  $X = \{\mathbf{r}_i, \mathbf{v}_i\}$ ,  $i = 1, \dots, N$ , can be considered as a set of  $N$  points in six dimensional one particle space.

Dividing up this one particle space into cells  $\Delta_\alpha$ ,  $\alpha = 1, \dots, n$ , centered on  $(\mathbf{r}_\alpha, \mathbf{v}_\alpha)$ , of volume  $|\Delta_\alpha|$ , we can describe the macro (meso) states of the system  $X \in \Gamma_E$  by specifying, with some leeway, the fraction of particles in each  $\Delta_\alpha$ . We shall denote by  $N_\alpha(X)$  the number of particles in  $\Delta_\alpha$ .

## Lanford's Theorem

Consider a gas consisting of  $N$  spheres of diameter  $d$  in a volume  $V \subset \mathbb{R}^3$  evolving according to Hamiltonian dynamics with elastic collisions.

Keeping  $V$  fixed consider now a sequence of systems with different particle numbers, and different diameters  $d$ , such that  $N \rightarrow \infty$ ,  $d \rightarrow 0$ , while  $Nd^2 \rightarrow b > 0$  (and so  $Nd^3 \rightarrow 0$ ). This is called the Boltzmann-Grad (BG) limit.  $b^{-1}$  is proportional to the mean-free-path between collisions.

Let  $f_0(\mathbf{x}, \mathbf{v}) \geq 0, \mathbf{x} \in V \subset \mathbb{R}^3, \mathbf{v} \in \mathbb{R}^3$ , be a smooth distribution of integral one in the one particle space.

Define now, for each  $N$ ,  $\Gamma_{M_{f_0}}$  to consist of all the phase points  $X_N$  such that  $X_N \in \Gamma_{M_{f_0}} \subset \Gamma_E$ , if

$$N_\alpha(X_N)/N \cong \int_{\Delta_\alpha} f_0(\mathbf{x}, \mathbf{v}) d\mathbf{x}d\mathbf{v}, \quad \text{for all } \alpha \quad (7)$$

Here  $\Gamma_E$  (and  $\Gamma_{M_{f_0}}$ ) will have  $6N$  dimensions but  $f_0(\mathbf{x}, \mathbf{v})$  is a fixed function on  $\mathbb{R}^6$ . As  $N$  increases the fraction of particles in  $\Delta_\alpha$ , is required to satisfy (7) ever more closely with equality in the BG limit,

$$\lim_{\text{BG}} N_\alpha(X_N)/N = \int_{\Delta_\alpha} f_0(\mathbf{x}, \mathbf{v}) d\mathbf{x}d\mathbf{v} \quad (8)$$

for arbitrarily small  $\Delta_\alpha$ .

Lanford's theorem then says:

Given  $f_0$ , a microstate  $X_N(0) \in \Gamma_{M_{f_0}}$  will typically evolve, via the Hamiltonian dynamics, to a microstate  $X_N(t)$  such that for all reasonable sets of  $\Delta_\alpha$ 's

$$\frac{N_\alpha(X_N(t))}{N} \cong \int_{\Delta_\alpha} f(\mathbf{x}, \mathbf{v}, t) d\mathbf{x}d\mathbf{v} \quad (9)$$

where  $f(\mathbf{x}, \mathbf{v}, t)$  solves the well-known Boltzmann equation

$$\frac{\partial f}{\partial t} + \mathbf{v} \cdot \frac{\partial f}{\partial \mathbf{x}} = Q(f, f) \quad (10)$$

with initial condition  $f_0(\mathbf{x}, \mathbf{v})$ . Eq. (9) means that so long as  $N$  is sufficiently large, the approximation will hold to an arbitrary degree of accuracy with arbitrary high probability with respect to the uniform measure on  $\Gamma_{M_{f_0}}$ .

The BE gives a deterministic evolution from the macrostate  $M_{f_0}$  to  $M_{f_t}$  in the BG limit. It will hold to a good approximation for a macroscopic dilute gas. It accurately describes the typical time evolution of the "empirical" distribution of atoms in the one particle space of such an individual system.

The Boltzmann entropy of a dilute gas in a macrostate  $M_f$  (as defined in (5)) is,

$$S_B(f) = S_B(M_f) = \log |\Gamma_{M_f}| \quad (11)$$

where  $|\Gamma_{M_f}|$  is for an  $N$  particle system the  $6N$  dimensional phase space volume of  $\Gamma_{M_f}$

$S_B(f)$  was actually computed by Boltzmann. He showed that, up to constants, this is given for a dilute gas of  $N$  particles,  $N \gg 1$ , by

$$\frac{1}{N} S_B(f) = - \int_V d\mathbf{x} \int_{\mathbb{R}^3} d\mathbf{v} f(\mathbf{x}, \mathbf{v}) \log f(\mathbf{x}, \mathbf{v}) + \text{Const.} \quad (12)$$

While (12) looks like, and is sometimes mistakenly confused with, the Gibbs-Shannon entropy per particle for a system in a product measure of the form  $\prod_i f(\mathbf{x}_i, \mathbf{v}_i)$ , it is conceptually not the same at all.  $S_B(f)$  is the entropy of an individual system with empirical distribution  $f(\mathbf{x}, \mathbf{v})$  (see below).

The maximum of  $S_B(f)$  over all  $f$  for a system of  $N$  particles in a volume  $|V|$  with a given energy, which is for a dilute gas just the kinetic energy, is given by the Maxwell distribution

$$f_{\text{eq}} = (2\pi kT/m)^{-3/2} \exp[-m\mathbf{v}^2/2kT]/|V| \quad (13)$$

where  $kT = 2/3(E/N)$ .

The distribution  $f_{\text{eq}}(\mathbf{x}, \mathbf{v})$  is the unique stationary solution of the BE.

In this case  $\Gamma_{M_{f_{\text{eq}}}} = \Gamma_{M_{\text{eq}}}$  and the entropy/particle is given by

$$\frac{1}{N} S_B(f_{\text{eq}}) = \frac{3}{2} \log T - \log \frac{N}{|V|} + \text{Const.} \quad (14)$$

the same as the equilibrium Clausius entropy for a dilute gas, with density  $\frac{N}{|V|}$ .

The second law now says that for a *typical* microstate of a dilute gas with  $X(0) \in \Gamma_{M_{f_0}}$ ,  $f_0 \neq f_{\text{eq}}$ ,  $f(\mathbf{x}, \mathbf{v}, t) = f_t$  has to be such that  $S_B(f_t) \geq S_B(f_{t'})$ , for  $t \geq t'$ .

This is exactly what happens for  $f(\mathbf{x}, \mathbf{v}, t)$  evolving according to the Boltzmann equation:

$$\frac{d}{dt} S_B(f_t) \geq 0, \quad \text{Boltzmann's } \mathcal{H}\text{-theorem} \quad (15)$$

As put by Boltzmann<sup>4</sup>:

*“In one respect we have even generalized the entropy principle here, in that we have been able to define the entropy in a gas that is not in a stationary state.”*

---

<sup>4</sup>Bo98

Lanford could prove his theorem only for a time  $0 < t < \tau$ . This is still the case at present. In fact one can only prove the existence of smooth solutions of the BE by using Lanford's theorem and then of course only for times  $t < \tau$ .

The time  $\tau$  for which Lanford's theorem holds is about one fifth of the mean free time between collisions, but that is a purely technical problem. This time is long enough for the Boltzmann entropy per particle to increase by a finite amount.

N.B. The collision term in the Boltzmann eq. (10) is proportional to  $b = \lim_{BG} Nd^2$ . If one takes the limit  $N \rightarrow \infty$ ,  $d \rightarrow 0$  in such a way that  $Nd^2 \rightarrow 0$  then the equation for  $f$  is just the same as for an ideal gas

$$\frac{\partial f}{\partial t} + \mathbf{v} \cdot \frac{\partial f}{\partial \mathbf{x}} = 0. \quad (16)$$

It follows then that the time derivative of  $S_B$  given in (12) is equal to zero.

The reason for this is that as  $b$  gets smaller the time between collisions gets larger and when  $b \rightarrow 0$  this time becomes infinite. The one particle distribution  $f(x, v, t)$ , given in (16) will then actually describe, for times in which no collisions (interactions) have taken place (any fixed  $t$  when  $b = 0$ ), the time evolution of the microstate of the system, as specified by the  $N$  points in the one particle space, for  $N \rightarrow \infty$ . It therefore does not capture the features of the evolution of macrostates, essential for the increase of the Boltzmann entropy, i.e. the increase of  $|\Gamma_M(t)|$ .

What happens to the Boltzmann entropy of an ideal gas when  $M$  is a genuine macroscopic description has been investigated recently by us numerically and analytically for a one dimensional ideal gas in a box<sup>5</sup>. We considered a fixed decomposition of the one particle phase space  $(x, v)$  into rectangles of sizes  $|\Delta_\alpha| = \Delta x \Delta v$ , and defined the macrostate  $M$ , by the  $\{f_\alpha\}$ ,  $f_\alpha = \frac{N_\alpha}{|\Delta_\alpha|}$  where  $N_\alpha$  is the number of particles in the box  $\alpha$ . We then have, for large  $N$ ,

$$S_B(M) = - \sum_{\alpha} |\Delta_\alpha| f_\alpha \log f_\alpha \quad (17)$$

up to a constant.

---

<sup>5</sup>S. Chakraborti, A. Dhar, S. Goldstein, A. Kundu and J. Lebowitz (arXiv:2109.07742)

The time evolution of  $S_B(M(X_t))$  when  $X_0$  is chosen at random from an initial  $\Gamma_{M_0}$  corresponding to the phase space region where all  $N$  particles are in in the left half of the box with energy  $E$  approximately  $N\epsilon_0$  is shown in Figure 2 where  $s_B^f(t) = \frac{1}{N}S_B(M(X_t))$ .




Figure 2: A comparison between  $s_B^f(t)$  obtained from simulation of a single realization, and  $s_\Delta^F(t)$  obtained from an ensemble average, plotted as a function of time during free expansion.

For evaluating  $s_B^f(t)$  we use numerical simulation of a single initial condition  $X(0)$  picked at random from  $\Gamma_{M_0}$  for  $N = 10^7$  in a box of size  $L = 4$  (arbitrary units). We compute  $s_B^f(t)$  for different grid sizes  $|\Delta_\alpha| = \Delta x \Delta v$  by keeping  $\Delta x = 0.5$  fixed and varying  $\Delta v = 0.5$  (red empty circles), 0.25 (blue empty inverted triangles), 0.1 (yellow empty circles), and 0.05 (green empty triangles).

The solid lines,  $s_\Delta^F(t)$ , are the Gibbs-Shanon entropies obtained analytically from a coarsegrained one particle density in the  $(x, v)$  space. That is we set

$$F(x, v, 0) = \sum_{i=1}^N \langle \delta(x - r_i) \delta(v - v_i) \rangle, \quad (18)$$


where the average is over the uniform microcanonical density for  $X = \{r_1, \dots, r_N, v_1, \dots, v_N\}$  in  $\Gamma_{M_0}$ . We then evolve  $F$  according to Eq. (16) and then coarsegrain over cells  $\Delta_\alpha$ .

The evolution of an ideal gas on the interval  $[0, L]$  can be mapped, see (CDGKL), into the evolution on a circle of circumference  $2L$  and for simplicity of notation we shall sometimes set  $2L = 1$ . This does not affect anything relevant, provided we replace  $F$  below by its images under the mapping.

The figure suggests that  $s_B^f(t) = s_B^f(0)$  when  $\Delta v \rightarrow 0$  for any fixed  $t$ . This is consistent with what we found for the case where  $Nd^2 \rightarrow 0$ .

Note however that for each fixed  $\Delta v$ , no matter how small,  $s_B^f(t)$  seems to go, as  $t \rightarrow \infty$ , to a final equilibrium value, which for the initial condition considered here is just  $s_B^f(0) + \log 2$  as expected from the second law.

In fact the different curves  $s_B^f(t)$  collapse to a single curve when  $t$  is rescaled  $t \sim \frac{\tau}{\Delta v}$ , see Figure.



**Figure 3:** The figure shows a collapse of the data presented in Figure 2 for different values of  $\Delta v$ , on plotting the entropy as a function of the scaled time  $\tau = t\Delta v/(2L)$ . Inset: A zoom up of the plot. The horizontal dotted line corresponds to the final equilibrium entropy.

The dashed line is the graph of  $s_B^F(\tau)$ , defined below for which it can be shown that

$$s_B^F(\tau) = \lim_{\Delta \rightarrow 0} s_B^F(2L\tau/\Delta v). \quad (19)$$

The quantity  $s^{\bar{F}}(\tau)$  is given by

$$s^{\bar{F}}(\tau) = -\frac{1}{N} \int \bar{F}(x, v, \tau) \ln \bar{F}(x, v, \tau) dx dv \quad (20)$$

where setting  $2L = 1$ , and considering the motion in a periodic box of unit length,

$$\bar{F}(x, v, \tau) = \frac{1}{\tau} \int_0^\tau F(x - x', v, 0) dx'. \quad (21)$$

## Setting

$$F(x, v, 0) = \rho h(v)[1 + \phi(x, v)], \quad (22)$$

$$\phi(x, v) = \phi(x + 1, v), \quad \int_0^1 \phi(x, v) dx = 0. \quad (23)$$

$$\bar{F}(x, v) = \rho h(v)[1 + \frac{1}{\tau} \int_0^\tau \phi(x - x', v) dx'] \quad (24)$$

$$= \rho h(v)[1 + \frac{1}{\tau} \psi(x, v, \tau)], \quad (25)$$

where  $\psi(x, v, \tau)$  is periodic in  $\tau$  with period 1 with

$$\psi(x, v, n) = 0, \quad \text{for } n = 1, 2, 3, \dots \quad (26)$$

This yields the  $s\bar{F}(\tau)$  plotted in Figure 3 for the initial condition used in Figure 2. The maxima, all of equal height occur at  $\tau = 1, 2, \dots$

## More General Macroscopic Equations

Going beyond the BE suppose, more generally, that the time evolution of the macrostate  $M$ , given by  $M(X(t)) = M_t$ , effectively satisfies an autonomous deterministic time asymmetric equation, such as the diffusion or the heat equation. (I shall consider here for simplicity macrostates  $M$  which are invariant under velocity reversal.)

Having an equation like the Boltzmann equation just discussed means that if  $t_3 > t_2 > t_1$ , then the microscopic dynamics  $T_t$  carries  $\Gamma_{M_{t_1}} = \Gamma_{M_1}$ , etc., inside  $\Gamma_{M_2}$  and  $\Gamma_{M_2}$  inside  $\Gamma_{M_3}$ , i.e.  $T_{t_2-t_1}\Gamma_{M_1} \subset \Gamma_{M_2}$  and  $T_{t_3-t_2}\Gamma_{M_2} \subset \Gamma_{M_3}$ , with *negligible error*. Put otherwise a typical phase point in  $\Gamma_{M_1}$  will go to  $\Gamma_{M_2}$  and then to  $\Gamma_{M_3}$ , i.e.  $T_{t_3-t_1}\Gamma_{M_1} \subset \Gamma_{M_3}$ .




Figure 4: Time evolution of  $\Gamma_{M_1}$

The fact that phase space volume is conserved by the Hamiltonian time evolution implies that  $|\Gamma_{M_1}| \leq |\Gamma_{M_2}| \leq |\Gamma_{M_3}|$ , and thus that  $S_B(M_3) \geq S_B(M_2) \geq S_B(M_1)$ .

Hence the solution of any deterministic macroscopic equation for  $M_t$  has to satisfy the inequality<sup>6</sup>  $\frac{d}{dt} S_B(M_t) \geq 0$ , exactly what Boltzmann showed for the BE.

---

<sup>6</sup>Go04,Pe70

We note that the existence of such a macroscopic equation implies (at least insofar as the macro-variables are concerned) that the phase points in the region in  $\Gamma_2$  coming from  $\Gamma_1$  behave, forward in time, as microstates typical of  $\Gamma_2$ .

They are, however, very atypical backwards in time. Thus if we reverse all the velocities in  $\Gamma_2$ , then at a later time,  $t' = t_2 + (t_2 - t_1)$  all of the points initially in  $\Gamma_{M_1}$  will again be in  $\Gamma_{M_1}$  (with their velocities reversed), a smaller region than  $\Gamma_{M_2}$ , while  $|\Gamma_{M_{t'}}| \geq |\Gamma_{M_2}|$ .

The reason for this asymmetry in typical behaviors is due to initial conditions. That is, when nature or the experimentalist who is part of nature, starts out at  $t_1$  with a nonequilibrium system in an initial state  $X \in \Gamma_{M_1}$  we can assume, for  $X$  typical of  $\Gamma_{M_1}$ , that  $X(t)$  will continue to be typical of  $M_t$  insofar as times greater than  $t_1$  are concerned but obviously not with regard to times earlier than  $t_1$ .

This was shown explicitly in the Lanford derivation of the BE: reversing the velocities at some  $t > 0$  ( $t < \tau$ ) violates the assumptions on the initial conditions required for the derivation of the Boltzmann equation.

If we could indeed **perfectly** reverse all the velocities in a macroscopic system out of equilibrium and keep the system **perfectly** isolated afterwards we would have a violation of the second law.

But what about real life situations?

What corresponds to an appropriate choice of initial time and initial conditions for which one can, in fact has to, assume that at that time the state  $X$  is atypical of  $\Gamma_E$  (because it is a low entropy nonequilibrium state)? Somewhat surprisingly, if one thinks hard about it, one is pushed to consider the very beginning of the universe we live in.

This would correspond according to our current physical theories to the time just after the “Big Bang”. The importance of initial conditions, Big Bang or not, was already fully understood by Boltzmann and others as the quotes below show.

## Initial Conditions

*“From the fact that the differential equations of mechanics are left unchanged by reversing the sign of time without changing anything else, Herr Ostwald concludes that the mechanical view of the world cannot explain why natural processes always run preferentially in a definite direction. **But such a view appears to me to overlook that mechanical events are determined not only by differential equations, but also by initial conditions.** In direct contrast to Herr Ostwald I have called it one of the most brilliant confirmations of the mechanical view of Nature that it provides an extraordinarily good picture of the dissipation of energy, as long as one assumes that the world began in an initial state satisfying certain conditions. I have called this state an improbable state.”*


— L. Boltzmann<sup>7</sup>

---

<sup>7</sup>Bo97

*“It is necessary to add to the physical laws the hypothesis that, in the past the universe was more ordered in the technical sense, [i.e. low  $S_B$ ] than it is today ... to make an understanding of irreversibility.”*

— R.P. Feynman<sup>8</sup>



**Figure 5:** “Creation of the universe: a fanciful description! The Creator’s pin has to find a tiny box, just 1 part in  $10^{10^{123}}$  of the entire phase-space volume, in order to create a universe with as special a Big Bang as we actually find.” from R. Penrose, *The Emperor’s New Mind*

The “tiny box” in the figure is a macrostate with low  $S_B$ . N.B. It is not necessary to select a particular microstate. Almost all microstates in a low-entropy macrostate will behave in a similar way.

It may be relevant to mention here a question I was asked during a talk I gave on the subject:

Q: What does the initial state of the universe have to do with the fact that when I put my sugar cube in my tea it dissolves irreversibly?

A: Nothing directly. But the fact that you, the sugar cube and the tea are all here is a consequence of the initial low entropy state of the universe.

## Boltzmann vs. Gibbs Entropies

Given an ensemble (probability) density  $\mu(X)$ ,  $X \in \Gamma$ , the Gibbs-Shannon entropy is given by

$$S_\mu \equiv - \int_\Gamma \mu \log \mu \, dX. \quad (27)$$

Clearly if  $\mu = \tilde{\mu}_M$ , where

$$\tilde{\mu}_M = \begin{cases} |\Gamma_M|^{-1}, & \text{if } X \in \Gamma_M; \\ 0, & \text{otherwise} \end{cases} \quad (28)$$

then

$$S_G(\tilde{\mu}_M) = \log |\Gamma_M| = S_B(M). \quad (29)$$

This is essentially the case for the microcanonical ensemble since  $\Gamma_E \simeq \Gamma_{M_{\text{eq}}}$ . By the equivalence of ensembles for macroscopic systems the same is true for the canonical and other Gibbs ensembles.

Thus the Gibbs and Boltzmann entropies are equal for equilibrium systems.

However, as  $\mu = \mu_t$  evolves via the Hamiltonian dynamics for isolated systems  $S_G(\mu)$  does not change in time.  $S_G(\mu)$  is therefore “useless” for such systems not in equilibrium, while  $S_B(M(X_t))$  captures the essence of typical macroscopic behavior. In particular it satisfies the second law of thermodynamics.



**Figure 6:** Boltzmann's grave in Zentralfriedhof, Vienna, with bust and entropy formula

## Summary of Boltzmann's Ideas (also Maxwell, Kelvin, ...)

Time-asymmetric behavior as embodied in the second law of thermodynamics is observed in *individual macroscopic* systems. It can be understood as arising naturally from time-symmetric microscopic laws when account is taken of a) the great disparity between microscopic and macroscopic sizes, b) initial conditions, and c) that what we observe are “typical” behaviors — not all imaginable ones. Common alternate explanations, such as those based on equating irreversible macroscopic behavior with ergodic or mixing properties of ensembles (probability distributions) already present for chaotic dynamical systems having only a few degrees of freedom or on the impossibility of having a truly isolated system, are either unnecessary, misguided or misleading.

Let me end this section on systems described by classical mechanics by quoting Einstein's tribute to Boltzmann.

*“On the basis of kinetic theory of gases Boltzmann had discovered that, aside from a constant factor, entropy is equivalent to the logarithm of the “probability” of the state under consideration. Through this insight he recognized the nature of course of events which, in the sense of thermodynamics, are “irreversible”. Seen from the molecular-mechanical point of view, however all courses of events are reversible. If one calls a molecular-theoretically defined state a microscopically described one, or, more briefly, micro-state, then an immensely large number ( $Z$ ) of states belong to a macroscopic condition.  $Z$  is then a measure of the probability of a chosen macro-state. **This idea appears to be of outstanding importance also because of the fact that its usefulness is not limited to microscopic description on the basis of mechanics.”***

— A. Einstein, Autobiographical notes

# Quantum Systems

Let me turn now to quantum systems. These are very different on the microscopic level from, but quite similar on the macroscopic level to, classical systems.

The first question is: what is the best quantum description of the microstate of a physically isolated macroscopic system? Unfortunately there is no consensus on this, even a century after the birth of quantum mechanics, and its perfect success in explaining and predicting microscopic phenomena.

There is however no doubt that, as put by Einstein, Boltzmann's ideas are applicable to the quantum (real) world.

I will assume for this talk that the microstate of a macroscopic system with an energy in a narrow interval  $(E, E + \Delta E)$  to be a wave function  $\psi \in \mathcal{H}_E$ , where  $\mathcal{H}_E$  is a shell in Hilbert space of thickness  $\Delta E$ , i.e.  $\mathcal{H}_E$  consists of all linear combination of energy eigenfunctions in the range  $(E, E + \Delta E)$ ,  $\Delta E \ll E$  but very large compared to the spacing between levels.

This is not perfect (problems with Schrödinger's Cat) but it will have to do for the present (see below). Using a density matrix with eigenstates in  $\mathcal{H}_E$  would give similar results.

The next question then is which  $\psi$  correspond to the system being in macroscopic thermal equilibrium (MATE), i.e. what is the analog of a classical microstate  $X$  lying in  $\Gamma_{M_{\text{eq}}}$ .

Following von Neumann<sup>9</sup>, we take the macro-observables corresponding to the macrostate  $M$  to commute with each other. We assume that this can be achieved by suitably “rounding off”, i.e. coarse-graining, the operators representing the macro-observables.

---

<sup>9</sup>Chapter 5 of Ne55

The coarse-grained energy operator commutes with the other coarse-grained macro-observables, which I shall also denote by  $M$ . Thus all  $M$ 's can be regarded as operators on  $\mathcal{H}_E$ . Their joint spectral decomposition defines an orthogonal decomposition

$$\mathcal{H}_E = \bigoplus_{\nu} \mathcal{H}_{\nu}, \quad (30)$$

Each subspace  $\mathcal{H}_{\nu}$  consists of all linear combinations of the joint eigenvectors of the macro-observables whose eigenvalues lie in a narrow range. These subspaces  $\mathcal{H}_{\nu}$  correspond to the different macrostates and the orthogonal decomposition (30) of  $\mathcal{H}_E$  corresponds to the division of the classical energy shell  $\Gamma_E$  into disjoint regions  $\Gamma_M$ .

A system is in a macrostate  $M_\nu$  if its wave function  $\psi$  is “close” to  $\mathcal{H}_\nu$ , i.e.

$$\langle \psi | P_\nu | \psi \rangle \geq 1 - \delta, \quad \delta \ll 1, \quad (31)$$

with  $P_\nu$  being the projection to  $\mathcal{H}_\nu$ .

As noted earlier, due to the Schrödinger’s Cat problem, there will be  $\psi$ ’s which are a superposition of  $\psi$ ’s in different macrostates.

To remedy this one has to go beyond the Copenhagen doctrine that the wave function is a complete description<sup>10</sup>. For the present let me say that I would interpret such  $\psi$ ’s when they arise physically as giving probabilities  $\langle \psi | P_\nu | \psi \rangle$  of being in different macrostates,  $M_\nu$ .

---

<sup>10</sup>Be87,Go98

The analog of the phase space volume  $|\Gamma_M|$  is the dimension  $d_\nu$  of each macro space  $\mathcal{H}_\nu$ .

As in the classical case, it is generally true that one of the  $\mathcal{H}_\nu$ , denoted  $\mathcal{H}_{\text{eq}}$ , has most of the dimensions of  $\mathcal{H}_E$ , i.e.,

$$\frac{\dim \mathcal{H}_{\text{eq}}}{\dim \mathcal{H}_E} = 1 - \varepsilon \quad (32)$$

with  $\varepsilon \ll 1$ .

The Boltzmann entropy  $S_B$  of a system in a macrostate  $M_\nu$  is given by the log of the dimension of the macro space  $\mathcal{H}_\nu$ ;

$$S_B(M_\nu) = \log d_\nu. \quad (33)$$

Given this correspondence between partitioning  $\Gamma_E$  and decomposing  $\mathcal{H}_E$  most everything we said about isolated classical macroscopic systems then also holds for quantum systems. In particular for  $\varepsilon \ll \delta$ , most  $\psi \in \mathcal{H}_E$  are in MATE.

The Boltzmann argument for increase in entropy of isolated macroscopic systems out of equilibrium is then similar to that in the classical case<sup>11</sup>.

When I say “most  $\psi$ ”, I mean with respect to a uniform measure on the unit sphere in  $\mathcal{H}_E$ . This measure on “wave functions” was considered already by Schrödinger and particularly by Felix Bloch<sup>12</sup>. It yields the microcanonical density matrix  $\rho^{\text{mc}}$  but goes beyond it, in ascribing probabilities to any subset of  $\mathcal{H}_E$ .

---

<sup>11</sup>Gr94

<sup>12</sup>Go06

There is however also something beyond MATE which can be used for the characterization of quantum systems in equilibrium.

Unlike classical systems, where any subsystem of a system in a microstate  $X$  is also in a microstate  $X_S$ , having definite positions and velocities, a subsystem  $S$  of a quantum system with a wave function  $\psi$  will usually not be described by a wave function, but rather by a density matrix  $\rho_S^\psi$ , where

$$\rho_S^\psi = \text{tr}_{S^c} |\psi\rangle\langle\psi| \quad (34)$$

is the reduced density matrix of  $S$  obtained by tracing out the complement  $S^c$  of  $S$ .

We can now define<sup>13</sup> a system with wave function  $\psi$  to be in microscopic thermal equilibrium (MITE) if, for any not-too-large subsystem  $S$ , say subsystems with linear dimension  $l < l_0$ , for an appropriate  $l_0$ , the reduced density matrix of  $S$  is close to the thermal equilibrium density matrix of  $S$

$$\rho_S^\psi \approx \rho_S^{\text{mc}}, \quad (35)$$

$$\rho_S^{\text{mc}} = \text{tr}_{S^c} \rho^{\text{mc}}, \quad (36)$$

where  $\rho^{\text{mc}}$  is the microcanonical density matrix corresponding to a uniform distribution over energy eigenstates in  $\mathcal{H}_E$ . For macroscopic systems  $\rho_S^{\text{mc}}$  can be replaced by  $\rho_S^{\text{ca}}$ , where  $\rho^{\text{ca}}$  is the canonical density matrix.

The distinction between MITE and MATE is important for systems with many-body localization (MBL) for which most, if not all, the energy eigenfunctions fail to be in MITE while necessarily most of them, but not all, are in MATE.

---

<sup>13</sup>Go17

The argument for most energy eigenfunctions being in MATE is based on the fact that, calling  $D$  the dimension of  $\mathcal{H}_E$ , we have for energy eigenfunctions  $|n\rangle$

$$\frac{1}{D} \sum_{n=1}^D \langle n | P_{\nu_{\text{eq}}} | n \rangle = \frac{1}{D} \text{tr}(P_{\nu_{\text{eq}}}) = 1 - \varepsilon \quad (37)$$

Noting that  $\langle n | P_{\nu_{\text{eq}}} | n \rangle \leq 1$ , the average being close to 1 means that most eigenstates are close to  $\mathcal{H}_{\nu_{\text{eq}}}$ . This is consistent with the Eigenstate Thermalization Hypothesis (ETH).

In fact for generic macroscopic systems, including those with MBL, most wave functions in an energy shell are in both MATE and MITE. This follows from the following result.

## Canonical Typicality

Consider an isolated system consisting of two parts. Call them system 1 and 2 or system and reservoir. Then<sup>14</sup> we have the following result.

Let  $H$  be the Hamiltonian of the whole system and let the number of particles in system 1 and 2 be  $N_1 \ll N_2$ . Let  $\mathcal{H}_E \subset \mathcal{H}_1 \otimes \mathcal{H}_2$  be an energy shell. Then for most  $\psi \in \mathcal{H}_E$  with  $\|\psi\| = 1$ ,

$$\mathrm{tr}_2 |\psi\rangle\langle\psi| \approx \mathrm{tr}_2 \rho^{\mathrm{mc}}, \quad (38)$$

where  $\rho^{\mathrm{mc}}$  is the microcanonical density matrix of the whole system at energy  $E$ , i.e. equal weight to all energy eigenstates in  $\mathcal{H}_E$ .

When the interaction between systems 1 and 2 is weak then, as is well known,  $\mathrm{tr}_2 \rho^{\mathrm{mc}} \approx \frac{1}{Z} e^{-\beta H_1}$  for  $\beta = \beta(E) = dS_{\mathrm{eq}}(E)/dE$ .

---

<sup>14</sup>Go06,Le08

The theorem says that most wave functions in the energy shell  $\mathcal{H}_E$  are both in MATE and in MITE. In fact for macroscopic systems one can show that MITE implies MATE.

The opposite is however not true.

This is particularly relevant when one considers energy eigenfunctions  $|n\rangle$ . While most energy eigenstates, including those for systems with MBL must, as shown, generally be in MATE, most energy eigenfunctions for systems with MBL are not in MITE.

Interestingly MITE has been found experimentally and computationally to hold for systems with Hilbert spaces with dimension as small as 100 or so, for which the motion of macrostates is not really appropriate.

This brings up the question of how to apply what I have discussed here to systems with a small number of degrees of freedom. When considering such systems it is relevant to remember that probabilities of  $\mathcal{O}(10^{-6})$  might not be so different in practice from probabilities of  $\mathcal{O}(10^{-20})$ .

I will not discuss this or the related issues of “stochastic thermodynamics” here.

If MATE-ETH holds strictly, i.e., if *all* energy eigenstates in  $\mathcal{H}_E$  are in MATE, then every state  $\psi \in \mathcal{H}_E$  will sooner or later reach MATE and spend most of the time in MATE in the long run. That is because, writing  $\overline{f(t)} = \lim_{T \rightarrow \infty} \frac{1}{T} \int_0^T f(t) dt$  for time averages,  $|n\rangle$  for the energy eigenstate with eigenvalue  $E_n$  and  $\psi_t = e^{-iHt}\psi$ ,

$$\overline{\langle \psi_t | P_{\text{eq}} | \psi_t \rangle} = \sum_{n, n'} \langle \psi | n \rangle \overline{e^{iE_n t} \langle n | P_{\text{eq}} | n' \rangle e^{-iE_{n'} t} \langle n' | \psi \rangle} \quad (39)$$

$$= \sum_n |\langle \psi | n \rangle|^2 \langle n | P_{\text{eq}} | n \rangle \geq \sum_n |\langle \psi | n \rangle|^2 (1 - \delta) \quad (40)$$

$$= 1 - \delta, \quad (41)$$

provided  $H$  is non-degenerate, i.e.,  $E_n \neq E_{n'}$  for  $n \neq n'$  (using  $\overline{e^{iEt}} = 1$  if  $E = 0$  and  $= 0$  otherwise).

A similar statement is true when there is degeneracy.

## References

- [Be87] J.S. Bell (1987). *Speakable and Unspeakable in Quantum Mechanics*, Cambridge University Press, Cambridge, England.
- [Bo97] L. Boltzmann (1897). On Zermelo's Paper "On the Mechanical Explanation of Irreversible Processes". *Annalen der Physik*, **60**, 392-398.
- [Bo98] L. Boltzmann (1898). *Vorlesungen über Gastheorie*. Leipzig: Barth (Part I, Part II 1898). English translation by S.G. Brush: *Lectures on Gas Theory*. Berkeley: University of California Press (1964).
- [Ce94] C. Cercignani, R. Illner, M. Pulvirenti (1994). *The Mathematical Theory of Dilute Gases*. Berlin: Springer-Verlag.

[Fe67] R.P. Feynman (1967). *The Character of Physical Law*. Chapter 5. MIT Press, Cambridge, Mass.

[Go04] S. Goldstein, J.L. Lebowitz (2004). On the (Boltzmann) entropy of nonequilibrium systems. *Physica D*, **193**, 53-66.

[Go06] S. Goldstein, J.L. Lebowitz, R. Tumulka, N. Zanghi (2006). Canonical typicality. *Physical Review Letters*, **96**, 050403.

[Go98] S. Goldstein (1998). Quantum theory without observers. *Physics Today*, Part One: March, 42-46. Part Two: April, 38-42.

[Go17] S. Goldstein, D.A. Huse, J.L. Lebowitz, R. Tumulka (2017). Macroscopic and microscopic thermal equilibrium. *Annalen der Physik*, **529**, 1600301.

[Gr94] R. Griffiths (1994). Statistical Irreversibility: Classical and Quantum. Pages 147-159 in J.J. Halliwell, J. Pérez-Mercader, and W.H. Zurek (editors): *Physical Origin of Time Asymmetry*. Cambridge University Press.

[La73] O.E. Lanford (1973). Entropy and Equilibrium States in Classical Mechanics. Pages 1-113 in A. Lenard (editor): *Statistical Mechanics and Mathematical Problems*, Lecture Notes in Physics vol. **2**. Berlin:Springer-Verlag.

[La76] O.E. Lanford (1976). On a derivation of the Boltzmann equation. *Astérisque*, **40**, 117-137. Reprinted in J.L. Lebowitz and E.W. Montroll: *Nonequilibrium Phenomena — The Boltzmann Equation*, North-Holland (1983).

[Le08] J.L. Lebowitz (2008). From Time-symmetric Microscopic Dynamics to Time-asymmetric Macroscopic Behavior: An Overview. Pages 63-88 in G. Gallavotti, W.L. Reiter, J. Yngvason (editors): *Boltzmann's Legacy*. Zürich: European Mathematical Society.

[Ne55] J. von Neumann (1955). *Mathematical Foundations of Quantum Mechanics*. Pages 398-416. Princeton University Press, Princeton. Translated from German ed. (1932: Springer, Berlin) by R.T. Beyer.

[Pe70] O. Penrose (1970). *Foundations of Statistical Mechanics*. Pergamon, Elmsford, N.Y. Reprinted by Dover (2005).

[Pe90] R. Penrose (1990). *The Emperor's New Mind*. Chapter 7. Oxford University Press.